


NUTRITIONAL ADVANCES IN PRODUCTION PERFORMANCE AND PRODUCT QUALITY OF POULTRY HUSBANDRY UNDER HEAT STRESS

Syed Rizwan ALI SHAH¹✉, and Ibrahim Sadi ÇETİNGÜL¹

Afyon Kocatepe University, Department of Animal Nutrition and Nutritional Diseases, P.O.Box 03200, Afyonkarahisar, Turkey

✉Email: rizwansyed604@gmail.com;  ORCID: 0000-0002-2752-4854

↳Supporting Information

ABSTRACT: The objective of this review is to explore the nutritional additives to combat heat stress in high production targeted fowls. Many diseases and stress dynamics are liable for heavy mortality nowadays. High production targets and heat stress initiate free radical injuries and gastrointestinal oxidative insults resulting in poor bioavailability and feed efficiency. Accordingly, the quality of the eggs and meat is compromised along with the bird's health. Dietary schemes and relevant feed formulation with the provision of vital elements like selenium, zinc, calcium, vitamin E, vitamin C, vitamin A, electrolytes, essential amino acids, and plant extracts can conflict with all sorts of stress in birds and improve the immune system. The occurrence of many contagious diseases, nervous ailments, and metabolic syndromes can be decreased via appropriate feeding routines. It will not only control the bird's health but also increase the quality and market value of the products and consumer satisfaction. Scientists have struggled to prevent immuno-depression, egg and meat quality impairments by dietary influences. Poultry welfare is facing many problems currently which should be properly coped with innovative nutritional maneuvers. This review anticipates illuminating the probable nutritional approaches to manage stress in poultry birds.

Keywords: Heat stress, Poultry, Electrolytes, Phytogetic, Product quality

REVIEW ARTICLE
 PII: S222877012200008-12
 Received: January 05, 2022
 Revised: March 16, 2022
 Accepted: March 20, 2022

INTRODUCTION

The poultry sector faces several stress dynamics including feather pecking, vaccination, molting, increased stocking density, and high temperature. Although sometimes stress seems to be harmless, continuing stress circumstances can undesirably upset the vigor and production of the birds. Poultry birds need more nutrient requirements and metabolic rate is higher as compared to ruminants (Akşit et al., 2006). Poultry birds are very prone to heat stress as compared to other farm animals. Commercial broilers produce more metabolic heat as compared to other indigenous pure genotypes (Kang and Shim, 2021). Under heat stress, feed intake, feed conversion ratio, productivity, meat quality, egg quality, and immune status are compromised. Heat stress is a primary menace that deleteriously changes bird performance (Liu et al., 2020). Poultry birds uphold their body temperature at 40.6–41.0°C regardless of the ecological high temperature because of the homoeothermic activity (Ryder et al., 2004). Nevertheless, birds, whose bodies are covered with plumages and raised for high productivity are extra profound to heat stress (Mahmoud et al., 2015). Broiler meat and eggs are the most cost-effective animal protein nowadays. Globally poultry farming will progressively be exaggerated as birds are unprotected to periodic and prolonged heat stress associated with global warming (Wankhade et al., 2020).

The optimal temperature for the best productivity of birds is 19–22°C for laying hens and 18–22°C for growing broilers (Shahzad et al., 2021). High temperature in poultry causes reduced fertility, growth and production, poor quality meat and egg production, and more mortality (Ranjan et al., 2019). Furthermore, genetically modified layers kept for high production generate more metabolic heat so they are more prone to heat stress (Holik, 2015).

For the period of initial summertime, the poultry husbandry faces considerable damages in production, and particularly pale, soft, exudative (PSE) meat. PSE meat has a low water-holding ability and meat consistency. Certainly, these fluctuations are strictly interrelated to biochemical variations of skeletal muscles due to heat stress (Song and King, 2015). Immunity (Siddiqui et al., 2021), production, and bird welfare are undesirably affected due to heat stress in birds (Iyasere et al., 2021). The major reason for production loss due to stress is less feed intake of the birds and for every 1°C after 30°C, almost 5% feed intake drops which results in essential nutrient deficiencies (Bilal et al., 2021). To cope up with this, the supplementation of vitamin C and E, chromium, zinc, and methionine, high protein and energy, phytogetic extracts can be operative in decreasing the deleterious effects of heat stress (Khan et al., 2014; Surai et al., 2019).

This review is considered to anticipate the importance of nutrition in responding to the negative effects of heat stress on poultry production. Maximum of the possibilities and remedies of heat stress and negative impacts are illustrated.

DISCUSSION

Feeding and drinking response during stress

Undoubtedly feed intake decreases significantly when housing temperature upsurges. In a study, it was determined that feed intake was reduced and water intake was increased in broiler chicks and this behavior was in response to high metabolic heat (Chowdhury and Furuse, 2013). As a thumb rule, birds can be nourished at midnight during hot weather. Nevertheless, to avoid high metabolic heat overlapping ecological heat, it is usually recommended not to feed at 1.00-4.00 pm. Feed conversion rate or feed efficiency of the birds is reported to increase during heat stress periods irrespective of the feed intake and crop filling (Sohail et al., 2012). In a study, it was concluded that if broilers eat more finely ground feed particles during stress time, little metabolic heat is produced as compared to coarse particles of maize and the birds adapt to the environment with panting (Santos et al., 2019).

A study was conducted on 3060 broilers kept in tropical, subtropical, and temperate areas. FCR and body weight at 42 days were considerably higher ($p \leq 0.05$) in the subtropical area during winter (Osti et al., 2017). Temporary feed restriction appears to improve the productivity and feed efficiency of the birds during the hot season. A study was conducted on one-week-old broiler chicks kept in open housing and it was seen that feed restriction for 3 hours along with vitamins and mineral supplementation reduced mortality and improved production and quality of the meat (Mohamed et al., 2019). Similarly in another study, feed restriction for 2 hours per day improved the growth performance in broiler chicks (Liew et al., 2003). Conversely, feed restriction for a longer period decreased considerably the body weight gain and feed efficiency of broilers but reduced mortality (Lin et al., 2006).

It was shown in a study conducted on various genetic lines of broilers kept in semi-intensive housing and the results showed that all chicks showed maximum feeding intake at early morning 7:00 and late evening 17:00 when the temperature is decreased (Gonçalves et al., 2017). Hens under heat stress walk very little drink more water and do more rest as compared to control (Mack et al., 2013). The increase in water intake under stress is a physiological response of birds to homeostat body temperature and dehydration. Similarly in a study betaine was used in broilers due to its osmoregulation property to decrease heat damage and increase water intake (Ratriyanto and Mosenthin, 2018). The most economical way to decrease stress effects on birds is to feed them in the early morning and late evening and this will enhance feed efficiency (Liverpool-Tasie et al., 2019). Birds eat more in day time normally and less at night time but to compensate for the stress-induced decrease in feed intake in the daytime, nighttime feeding can be encouraged with synthetic illumination (Daghir, 2009).

Negative impacts of heat stress on poultry

Due to oxidative stress and cellular injuries during heat stress immunity and feed efficiency of the birds drops. There happens massive economic loss due to less production and mortality. The negative effects of heat stress on poultry are shown in figure 1 and table 1.

Nutritional intermediations improving product quality and production

Nutritional management is an auspicious way to enhance productivity, feed efficiency and reduce disease outbreaks by increasing the immunity of the birds during stress (Mujahid, 2011). Nutritional strategies include balancing of energy, provision of essential amino acids, adequate minerals and vitamins and many phytochemicals that are discussed below and shown in figure 2.




Figure 1 - Negative impacts of heat stress

Table 1 – The negative impacts of heat stress

Heat Stress	Effects	Birds	References
Chronic heat stress (10-days)	↓ Weight gain ↑ Inflammatory cytokines (IL-2 and TNF- α), and rectal Temperatures. ↑ Intestinal bacterial permeability	Broilers	Athenaky et al. (2017)
Chronic heat stress	↓ Feed intake and body weight.	Cobb-500 broiler	Attia et al. (2017)
Acute heat stress	↑ Mitochondrial Super-Oxide formation in bird's skeletal muscles.	Broilers	Sahin and smith (2016)
38°C (2, 5, and 10 hr.)	An efficient ↑ TLR4 expression in kidneys, liver, and heart.	28 days old broilers	Huang (2017)
23.9-37°C Cycling	↑ Heterophil-lymphocyte ratio, serum levels of cholesterol, glucose, and triglycerides. ↓ Feed intake, body and organ weight, primary and secondary antibody and IgM and IgG titers, HDL-cholesterol (serum concentration).	Broilers	Habibian et al. (2014)
31°C (12 HR.)	↓ In laying rate and egg production. ↑ (In mRNA levels) of ghrelin and amphetamine- and cocaine-regulated	12 weeks old hens	Song et al., (2012)
30°C (2 weeks)	↑ Serum corticosterone conc., and sodium-dependent glucose transporter-1 expression. ↓ Bodyweight.	28 days old broilers	Garriga et al. (2006)
32°C (acute heat stress)	Acid-base level disturbance (in the blood), negative effects on the integrity of muscle membrane, and changes in meat characteristics.	35-63 days old broilers	Sandercock et al. (2001)




Figure 2 - Nutritional strategies for decreasing heat stress

High-energy formulations

The benefits of giving high-energy formulations by supplementing fat and protein to broilers kept in high-temperature areas are familiar globally (Srinivasa Rao et al., 2016). For instance, a study was conducted on Ross-308 and chicks were fed with 21% and 23% crude proteins. The results showed a significant increase ($P < 0.05$) in body weight gain and feed conversion ratio as compared to the low protein diet. In short, it was declared that 23% protein and 3000 kcal/kg energy was better for growing stage and for the finishing period 20% protein and 3200 kcal/kg energy was suitable (Ahmed and Arabi, 2015). Feeding broilers with less protein-based diets during finishing has reduced microbial contents in feces and improved productivity during stress (Laudadio et al., 2012). In another study, for layers kept at 31 °C, supplementation of 5% fats to layers increased their feed intake by 17% while for hens kept at 11-18 °C, there was a 4.5% increase in feed intake (Daghir, 2008). Abdel-Moneim et al. (2020) reported that full-fat canola meal in Japanese quails under heat stress has resulted in improved growth and egg-laying along with egg quality.

Lowering metabolic heat

As compared to carbohydrates and protein, supplementary fat leads to a slight rise in metabolic heat (Laganá et al., 2007). Serving wet feed to broilers during stressful environmental conditions increases weight gain, feed conversion rate, and feed intake (Koh and Macleod, 1999). Mashed feed or crumbled feed particles are easily digested with little effort of gizzard i.e. high digestibility and hence reduced metabolic heat is produced (Amerah et al., 2007).

Betaine and methyl donors' supplementation

Betaine has good osmoregulation properties and its supplementation to quails at 0.06-0.12% has shown more feed intake and better egg quality during heat stress (Ratriyanto et al., 2017). Similarly in another study on quails, 16.5, 18, and 19.5% protein levels were augmented with 0, 0.06, 0.12% betaine respectively. The results showed that 18% protein with 0.12% betaine showed maximum crude fiber digestibility but less fat digestibility. Betaine improved all egg quality parameters and improved the digestibility of all nutrients (Raei et al., 2020). In another study, betaine was used in broiler chicks to decrease harmful effects of heat stress and it resulted in decreased mortality, improved body weight gain and feed intake, and humoral immune response of the chicks under stress in treatment groups as compared to the control (Ghasemi and Nari, 2020).

Reduced use of usual amino acids in diets during heat stress with supplementation of methionine, lysine, and betaine can improve the feed intake and efficiency along with a reduction in metabolic heat (Cronje, 2018; He et al., 2021). Moreover, these methyl group donors have value-added the carcass traits of broilers during heat stress (Ratriyanto et al., 2014). Glutamine supplementation in broilers under heat stress has resulted in intestinal epithelial cells multiplication increasing intestinal barrier function and tight junction proteins (Wu et al., 2018).

Taurine supplementation

Taurine has now made its position in animal nutrition as an antioxidant agent because of stress in poultry birds. In a study, taurine was used in broilers in drinking water for 35 days during heat stress. The results showed that there was a significant increase in weight gain and feed intake in taurine supplemented groups as compared to control. The gut fat was reduced along with improvement in muscle histopathological characters (Hafeez et al., 2021).

Probiotics and prebiotics supplementation

Probiotics and prebiotics are very valuable in poultry farming under heat stress. In laying hens provided with a high-temperature environment, 1g of mannan-oligosaccharides was used and it resulted in improvement of weight gain and feed conversion ratio, eggshell weight and thickness, albumen weight, and height in treatment groups (Bozkurt et al., 2012). Similarly, in broilers under heat stress, 24mg/kg of the diet was used for 42 days and it resulted in improvement of intestinal morphometric characters including villus width and surface area, and crypt depth and production of the birds (sohail et al. 2012). In laying hens probiotic mixture including 0.1% symbiotic and 0.1% culture of *Bacillus subtilis* and *Enterococcus faecium* and 0.5% mannan-oligosaccharide was used and it resulted in daily feed intake increase along with an increase in egg weight, egg-laying capacity, and egg quality parameters under heat stress. It also enhanced beneficial intestinal microbes (Zhang et al., 2017). In commercial broilers, strains of *Bacillus subtilis* were used and it increased daily body weight gain and lipid peroxidation reduction (Chao et al., 2017). Similarly, dietary supplementation of *Bacillus subtilis* at 0, 200 and 400 mg/kg of feed in broilers under heat stress increased significantly IgM levels in serum and improved cell-mediated immunity (Fathi et al., 2017). In a study, microalgae astaxanthin obtained from *Haematococcus pluvialis* was used in broilers and layers to evaluate its effects on molecular parameters of stress and lipid metabolism during heat stress and it decreased hepatic mRNA activity of many redox regulatory genes, heat shock transcription factor 1, heat shock protein 70 and tumor necrosis factor- α (Tolba et al., 2020). In a study, post-biotic obtained from *Lactiplantibacillus plantarum* was used in broilers to decrease the negative impacts of heat stress on production and bird's health. It resulted in upregulation of the glutathione peroxidase, superoxide dismutase, and catalase activity so it is a natural source of anti-oxidation (Humam et al., 2021).

Electrolytes supplementation

Electrolytes are very important for poultry that should be balanced during heat stress and it is recommended to be 250 mEq/kg during stressful conditions (Sarwar, 2007). For better productivity of broilers during heat stress, 0.21-0.25% sodium and 0.30% chloride should be provided during the finishing stage (Mushtaq et al., 2007). In a meta-analysis, it was predicted that for every 1°C increase in ambient temperature there was a 1.4% reduction in average daily feed intake and 2.1% reduction in average daily weight gain and dietary electrolyte balance has a strong effect on chicks less than 21 days old (Andretta et al., 2021). Due to decreased feed intake during heat stress, there comes dietary deficiency of vital minerals that are important for digestibility and oxidative stress neutralization. Hence, there should be controlled supplementation of calcium, zinc, iron, phosphorous, sodium, chloride, potassium, copper, magnesium, selenium, manganese, chromium, and iodine (Hassan Mir et al., 2018). In a study, broilers during the finishing stage were provided with minerals in water and feed during heat stress maintaining the electrolyte balance as 240 mEq. The results showed a 34% increase in water consumption as compared to control. Moreover, mineral supplementation reduced significantly metabolic heat, hyperventilation, and mortality (Farfán et al., 2010).

Vitamin C and vitamin E supplementation

Birds open to high-temperature present fall in the size of immune organs like the bursa, thymus, and spleen. Oxidative damage happens in birds prone to heat stress so, dietary addition of antioxidants is common dietary involvement. The provision of 250 mg/kg of vitamin E has been seen to decrease undesirable effects that occur during heat stress (Kazim Sahin et al., 2001). In another study, vitamin E was supplemented as 250 mg/kg of the diet in broilers and it resulted in increased activity of glutathione peroxidase and reduced expression of HSP60 (Kumbhar et al., 2018), and in layers, at 150 mg/kg it has shown to increase egg number and quality (Zhao et al., 2011). Ascorbic acid or vitamin C is being synthesized by the kidneys of the chicken and during heat stress, it becomes insufficient to cope with the bird's requirements. Vitamin C improves immunity, body weight, anti-oxidative potential, semen quality, and carcass characteristics in chicks (Abidin and Khatoun, 2013). A study was conducted to evaluate the effects of vitamin and mineral premix and vitamin C in broiler chicks during 0-35 days of the growing stage under artificially induced heat stress. The results showed that vitamin and mineral premix in water increased body weight gain and feed intake linearly, and the corticosterone level was reduced in serum in all treatment groups as compared to control (Saiz del Barrio et al., 2020). Similarly in another study, Vitamin C was provided to broiler chicks during the growing stage and the feed intake and body weight gain was improved due to a decrease in metabolic heat in birds as compared to the control group where no vitamin C was provided (Abudabos et al., 2018). In another study, Vitamin A, C, E, capsaicin from red pepper, selenium, and zinc were evaluated in layers kept under a heat stress environment. The supplementation increased weight gain, feed efficiency, and immune organs such as spleen and thymus weights, egg weight, shell thickness, and egg-laying percentage as compared to the control. Red pepper improved egg yolk color and decreased total cholesterol and lipids vice versa to selenium (Malekizadeh et al., 2013).

ALA and GAA supplementation

The α -Lipoic acid (ALA) and guanidinoacetic acid (GAA) are naturally synthesized in body and they are also present in many feed ingredients good antioxidants and if supplemented during heat stress, show positive response on production. In a study, α -Lipoic acid was used in broiler chicks kept at high temperatures and it increased average daily weight gain, feed intake, and volatile fatty acids in treatment groups as compared to the control. Moreover, villus height and villus to crypt depth ratio were improved along with the increase of beneficial microbes in the cecum (Wasti et al., 2021). In laying quails, guanidinoacetic acid (GAA) was supplemented during heat stress conditions and it resulted in increased laying capacity, better egg quality parameters, increased activity of superoxide dismutase, glutathione peroxidase, and reduction of malondialdehyde in serum (Raei et al., 2020).

Chitosan and galacto- oligosaccharides supplementation

In a study, chitosan oligosaccharides at 200 mg/kg were supplemented to broilers kept at high temperatures and it resulted that in the treatment group average daily weight gain, feed efficiency, weights of the immune organs, muscle oxidative enzyme activities were increased along with decreased malondialdehyde and corticosterone content as compared to control (Chang et al., 2020). Galacto-oligosaccharides were used in broilers kept at high temperatures and it resulted in improvement of feed conversion rate and weight gain along with a 5% reduction in mortality in treatment groups as compared to the control group (Slawinska et al., 2020).

Copper and zinc nanoparticles supplementation

In a study, nanoparticles of copper oxide were used to decrease heat stress effects on broiler chicks. The results showed that there was a reduction in metabolic heat and an increment in body weight gains in copper nanoparticles supplemented groups. Moreover, the stress biomarker, malondialdehyde (MDA) was decreased along with increased activities of catalase (CAT), superoxide dismutase (SOD), and glutathione peroxidase. Moreover, the liver tissue histology parameters were improved along with modulation of heat shock protein 70 and heat shock protein 90 (El-Kassas et al., 2019). In another study nanoparticles of zinc oxide were fed to layers as 80 mg/kg under heat stress and it resulted in a significant increase in egg numbers, egg weight, eggshell thickness, superoxide dismutase (SOD) activity, and zinc bioavailability in the treatment group as compared to control (Abedini et al., 2018).

Zinc, chromium, selenium, and methionine chelates supplementation

Effects of supplementation of zinc, chromium, and selenium in feeds were recorded in broiler chicks during the growing stage under heat stress. The results showed that there was an increase in feed intake and body weight gain in Zn, Se and Cr supplemented groups along with a reduction in lipid peroxidation as compared to the control. Moreover, the activity of superoxide dismutase was increased (Rao et al., 2016). Similarly, Zn bound to methionine and zinc oxide was supplemented to broilers under heat stress and it increased weight gain, nutrient bioavailability, thymus and bursa weights, egg quality, and reduction of malondialdehyde (MDA) and cholesterol in serum (Abd El-Hack et al., 2018). Similarly in another study, Zn-methionine (20, 40, and 80 mg/ kg of feed) provision to layers under heat stress has resulted in increased production, immunity, and egg quality parameters and reduced lipid peroxidation as compared to inorganic Zn and control group (Min et al., 2018). Chromium is well-thought-out to be a valuable trace element. It has been stated that its supplementation encourages the productivity of broilers during heat stress. In a study, chromium was

supplemented methionine in broilers and the results showed an increase in body weight gain, feed conversion ratio along with the coefficient of energy metabolism, serum glucose, cholesterol, and triacylglycerol (Santos Dalólio et al., 2021). Amino acid-bonded trace minerals were provided to broilers under heat stress and it resulted in a decrease of the corticosterone levels and cytokines in serum and resultantly improved productivity of chicks (Baxter et al., 2020). In another study, ethoxyquin and mineral-methionine chelate were used in broilers and it resulted in a reduction of development of wooden breasts in broilers and the lipid peroxidation of muscles was significantly lower as compared to control when both were used together (Kuttappan et al., 2021).

NaHCO₃ and KCL supplementation

Dietary supplementation of NaHCO₃ was performed in layers and immune response was improved against ND virus after vaccination, and digestibility of nutrients was significantly higher in treatment groups as compared to the control (Abbas et al., 2019). Potassium chloride was used in broiler chicks under heat stress and it increased the digestibility coefficients of dry matter, protein, fat, fiber, and crude ash. Moreover, the bioavailability of calcium and phosphorous was more for potassium chloride supplemented chicks as compared to the control. The red blood cells and total hematocrit were more in treatment groups as compared to the control (Gorlov et al., 2020).

Propolis supplementation

Propolis is obtained from honeybees and it has several beneficial effects under stressful environments. In a study on quails kept under high temperatures, propolis was used for minimizing the adverse effects of heat stress. It resulted in improvement of weight gain and feed intake, increases in intestinal crypts depth, and immune organ weights (Mehaisen et al., 2017).

Phytogenic supplementation

Many phytogenic feed supplements are being used nowadays to combat stress in poultry birds. Phytogenic feed additives have medicinal properties and natural to use without any side effects. They also enhance feed palatability with the flavor, taste and texture that the birds love to eat. Phytogenics enhance feed intake, promote gastrointestinal health status of the birds, improve immune system, reduce cellular oxidations, and improve production and product quality. There are many phytogenics that are mentioned below.

Ginseng

In a study, ginseng was used in broilers and it resulted in lowering the gene expression of HSPA1A, HSP1, and HSP-16.2, and it was reported to be the most suitable additive to reduce heat stress undesired effects (Sandner et al., 2020).

Dried plums

Dried plums were used in a study on broilers to reduce the harmful effects of heat stress. It increased the daily weight gain, feed intake, and feed conversion rate of the broilers, and it increased the expression of the genes related to anti-oxidation, heat shock proteins, and immune system activation genes as compared to the control (Wasti et al., 2021).

Watermelon juice

Watermelon juice was added as 20% and 40% to water and it was used during the finishing stage of the broilers for 21 days under heat stress. The results showed that the survival rate was improved and it was 100% in the 40% supplemented group. Moreover, weight gain and feed intake were increased significantly in all treatment groups as compared to the control (Jimoh et al., 2020).

Curcumin

Curcumin is obtained from turmeric and it was used in laying hens to improve the immunity during heat stress and it resulted in a reduction of Interleukins-6, interleukins 1 β , and tumor necrotic factor- α in treatment groups as compared to the control (Nawab et al., 2019). In another study, curcumin supplementation in broilers at 100 mg/kg of the diet (Zhang et al., 2018) and laying hens at 150 mg/kg of the diet (Liu et al., 2020) has increased growth performance and anti-oxidative and immune regulating gene activities, and egg and meat quality. In another study, curcumin nanoparticles were used in broilers and they were provided with 25, 50, and 100 mg/kg of the diet under heat stress. The results showed a significant increase in body weight gain, feed intake, immune organ weights reduction in malondialdehyde, and an increase in glutathione peroxidase and superoxide dismutase activity. Curcumin nanoparticles at 50 and 100 mg/kg of diet showed more significant results as compared to other groups (Badran, 2020). Similarly, cumin seed oil was supplemented to layers as 500 g/ ton of feed and it resulted in increased egg quality, feed conversion, immunity, and drop in serum malondialdehyde level (Saleh et al., 2019).

Cumin seeds

Cumin seed powder supplementation in broilers under heat stress has resulted in increased weight gain, feed conversion ratio and dressing percentage (Chawke et al., 2021). In a study, the effects of phytogenic feed additive

“comfort” (PFA-C) was evaluated in broilers and it resulted in down-regulation of heat shock protein 70 gene expression, increase in water and feed intake, and average weight gain and low mortality (Greene et al., 2021).

Cinnamon and clove

In a study, cinnamon and clove powders and their oils were used in quails reared under heat stress and it resulted in a significant drop in heat shock protein HSP70, HSP40, and HSP90, and tissue malondialdehyde, corticosterone, and cytokines in serum were reduced along with the increased activity of catalase, superoxide dismutase and glutathione peroxidase in treatment groups as compared to the control. Moreover, the oils were more beneficial as compared to the powders (Mustafa and Wasman, 2020). Similarly in broilers, cinnamon powder as 0.5% has increased the weight gain and activity of serum glutathione peroxidase, catalase, and superoxide dismutase enzymes and reduction in malondialdehyde content (Sadeghi and Moghaddam, 2018).

Slam weed

Siam weed (*Chromolaena odorata*) was used in broiler chicks to decrease the heat stress effects. Its Ethanol extract at 25–400 µg/ml presented a resilient antioxidant activity in vitro, comparable to 10–80 µg/ml ascorbic acid supplementation. Chromomoric acid obtained from siam weed (*Chromolaena odorata*) at 10 µg has anti-inflammatory activity. Moreover, the toxicity trials have shown that only the aerial fragments of it are nontoxic for animal feeding (Lartey et al., 2020).

Fennel seeds

A study was performed in laying hens where fennel (*Foeniculum vulgare*) at 0, 10, and 20 g/kg were used to decrease harmful effects of heat stress and it resulted that malondialdehyde and egg carbonyl contents were reduced along with egg triglyceride and yolk cholesterol contents (Gharaghani et al., 2015).

Herbal mixture

In a study, extracts of *Phyllanthus emblica*, *Tribulus terrestris*, *Withania somnifera*, *Ocimum sanctum* *Asparagus racemosus*, *Mangifera indica*, and *Glycyrrhiza glabra* were mixed and used in broilers under heat stress and it resulted in a significant increase in body weight and feed intake. Moreover, free radical formation, blood urea nitrogen, serum cholesterol, and glucose level were decreased significantly in treatment groups as compared to control at 42 days of age (Rao et al., 2021).

African moringa

African moringa (*Moringa stenopetala*) leaf meal was used in broilers and layers kept in a high-temperature environment and it resulted that 2% supplementation of it improved egg yolk color and there was no significant effect on egg and meat quality (Tamiru et al., 2020).

Lycopene

Lycopene is a carotenoid that is being found in tomatoes (*Solanum lycopersicum*) and it has shown anti-oxidative, anti-inflammatory activities along with improved productivity and feed efficiency, and better broiler meat and egg quality (Arain et al., 2017). Similarly in another study in laying hens, lycopene as 20 mg/kg of the diet has shown a decrease in lipid peroxidation and increase in egg yolk color along with mineral and vitamin levels (An et al., 2019).

Licorice

In a study, Licorice (*Glycyrrhiza glabra*) was used in egg-laying quails under heat stress and it resulted in a reduction of egg yolk cholesterol and total lipids in licorice fed groups at 500 mg as compared to the control. Moreover, total serum cholesterol, low-density lipoproteins, triglycerides, and total serum lipids were decreased significantly in treatment groups as compared to control (Dosoky et al., 2021).

Willow

Willow (*Salix*) bark has been used in poultry birds to alleviate the negative impact of heat stress on birds' health and production because of the important source of flavonoids, tannins, salicin, and glucosides of saligenin (Saracila et al., 2021). Organic acids and essential oils are safe and have been approved by European Union for supplementation in animal feeding as anti-oxidative, antimicrobial, immune stimulators, and stress relieving additives (Doneria et al., 2020).

Syrian oregano

In a study, Syrian oregano (*Origanum syriacum*) oil and Avilamycin, an antibiotic, effects were compared in broilers under heat stress for 42 days and the results showed no significant difference between the effects of the 600 mg/kg *Origanum syriacum* and avilamycin as compared to the control and it has shown improvement in production and immune status of the birds under heat stress (Tekce et al., 2021).

Garlic

Individual and combined effects of probiotic, garlic powder, citric acid was evaluated in broiler chicks under chronic heat stress. It resulted in improvement of productivity and feed efficiency, immunity, intestinal microbial population, and intestinal morphometric characters in treatment groups as compared to the control (Elbaz et al., 2021).

Thyme

In a study on rabbits, thyme leave powder was used for 90 days to minimize the deleterious effects of heat stress and it resulted in a significant increase in the feed intake, body weight gain, liver and kidney functionality, improved testosterone and semen parameters in all treatment groups as compared to the control and particularly for 16 g/kg diet supplementation of the thyme leave powder the results were highest for all the above parameters (Ahmed et al., 2020). In another study, thyme oil as 150, 200 mg/kg of diet in broilers under heat stress has resulted in a significant increase in weight gains, immunity, and reduced oxidative stress biomarkers (Khafar et al., 2019).

Onion

Onion powder as 2.5 kg/ton of feed and 2.5% onion extract in drinking water was used for broiler chicks under heat stress for 42 days and the results showed an increase in production and feed intake of the chicks, immunity, and heat tolerance as compared to control (Al-Ramamneh, 2018).

Grade seeds

In a study on the broiler, grape seed extract and vitamin C was supplemented for 42 days to cope with chronic heat stress and it resulted that 300 mg/kg grape seed extract caused a significant increase in body weight gain, decrease glucose, cholesterol, triglycerides and low-density lipoproteins in serum of the chicks of treatment groups as compared to the control (Hajati et al., 2015).

Saffron

Saffron (*Crocus sativus*) petal extract 300, 500, and 700 mg/kg of diet was used in broiler chicks for 42 days under heat stress and it resulted in a significant increase in feed intake and body weight gain, glutathione peroxidase, and superoxide dismutase activities and significant decrease of serum cholesterol, uric acid and malondialdehyde in treatment groups as compared to the control (Hosseini-Vashan and Piray, 2021).

Sweet wormwood

Sweet wormwood (*Artemisia annua*) was used in a study on broilers under heat stress to evaluate its antioxidant potential and relative meat quality parameters and it resulted in a significant increase of redness and energy status of breast muscles and decreased drip loss, reactive oxygen species production and the recommended level of use was 1-1.25 g/kg of diet (Wan et al., 2018).

Tulsi leaf and fenugreek seed

In a study, tulsi (*Ocimum sanctum*) leaf powder and fenugreek (*Trigonella foenum-graecum*) seed powder was used in broilers and the results showed that under a high level of temperature during experimentation the feed cost was not significantly changed but the production was improved in treatment groups as compared to the control (Prajapat et al., 2020).

Resveratrol (Red grapes)

Resveratrol obtained from red grapes was supplemented to layers under heat stress at 0, 200, 400, and 600 mg/kg of the feed and it resulted in improved productivity and humoral immunity and reduced serum malondialdehyde and gene expression of heat shock proteins (Liu et al., 2014). Similarly, ducks were supplemented with resveratrol under acute heat stress exposure and it resulted in increased villus height to crypt depth ratio, a higher number of goblet cells, increased tight junction proteins and reduced gastrointestinal artifacts, reduced gene expression of heat shock proteins (Yang et al., 2021). Similarly in another study on broilers, resveratrol supplementation has shown an increase in activity of superoxide dismutase and glutathione peroxidase and reduction in serum and meat malondialdehyde (He et al., 2019).

Green tea

Catechin extracted from green tea was supplemented to broilers under heat stress as 300 and 600 mg/kg of the feed and it resulted in a significant increase in weight gain and feed intake, enzymatic activities and immune response, and reduction in serum cholesterol as compared to control (Xue et al., 2017). Similarly in quails under heat stress catechin supplementation has resulted in increased feed intake and egg numbers and better antioxidant profile (Sahin et al., 2010).

Black cumin seeds

Black seeds (*Nigella sativa*) supplementation as 1-2% in broilers has increased weight gain and meat quality, reduced oxidation, and serum corticosterone levels (El-Shoukary et al., 2014).

- An BK, Choo WD, Kang CW, Lee J, and Lee KW (2019). Effects of dietary lycopene or tomato paste on laying performance and serum lipids in laying hens and on malondialdehyde content in egg yolk upon storage. *Journal of Poultry Science*, 56 (1): 52-57. [Article link](https://doi.org/10.2141/jpsa.0170118) | DOI: <https://doi.org/10.2141/jpsa.0170118>
- Andretta I, Kipper M, Schirmann GD, Franceschina CS, and Ribeiro AML (2021). Modeling the performance of broilers under heat stress. *Poultry Science*, 100 (9): 101338. [Article link](https://doi.org/10.1016/j.psj.2021.101338) | DOI: <https://doi.org/10.1016/j.psj.2021.101338>
- Arain M.A, Mei Z, Hassan F.U, Saeed M, Alagawany M, Shar A.H, and Rajput IR (2017). Lycopene: A natural antioxidant for prevention of heat-induced oxidative stress in poultry. *World's Poultry Science Journal*, 74 (1): 89-100. [Article link](https://doi.org/10.1017/S0043933917001040) | DOI: <https://doi.org/10.1017/S0043933917001040>
- Badran A (2020). Effect of dietary curcumin and curcumin nanoparticles supplementation on growth performance, immune response and antioxidant of broilers chickens. *Egyptian Poultry Science Journal*, 40 (1): 325-343. [Article link](https://doi.org/10.21608/epsj.2020.81756) | DOI: <https://doi.org/10.21608/epsj.2020.81756>
- Baxter MFA, Greene ES, Kidd MT, Tellez-Isaias G, Orlowski S, and Dridi S (2020). Water amino acid-chelated trace mineral supplementation decreases circulating and intestinal HSP70 and proinflammatory cytokine gene expression in heat-stressed broiler chickens. *Journal of Animal Science*, 98 (3): skaa049. [Article link](https://doi.org/10.1093/JAS/SKAA049) | DOI: <https://doi.org/10.1093/JAS/SKAA049>
- Bilal R.M, Hassan F.ul, Farag M.R, Nasir T.A, Ragni M, Mahgoub H.A.M, and Alagawany M (2021). Thermal stress and high stocking densities in poultry farms: Potential effects and mitigation strategies. *Journal of Thermal Biology*, 99:102944. [Article link](https://doi.org/10.1016/j.jtherbio.2021.102944) | DOI: <https://doi.org/10.1016/j.jtherbio.2021.102944>
- Bozkurt M, Küçükilymaz K, Çatli A.U, Çınar M, Bintaş E, and Çöven F (2012). Performance, egg quality, and immune response of laying hens fed diets supplemented with mannan-Oligosaccharide or an essential oil mixture under moderate and hot environmental conditions. *Poultry Science*, 91 (6): 1379-1386. [Article link](https://doi.org/10.3382/ps.2011-02023) | DOI: <https://doi.org/10.3382/ps.2011-02023>
- Chang Q, Lu Y, and Lan R (2020). Chitosan oligosaccharide as an effective feed additive to maintain growth performance, meat quality, muscle glycolytic metabolism, and oxidative status in yellow-feather broilers under heat stress. *Poultry Science*, 99 (10), 4824-4831. [Article link](https://doi.org/10.1016/j.psj.2020.06.071) | DOI: <https://doi.org/10.1016/j.psj.2020.06.071>
- Chao Y, Kim HW, Cramer T, Cheng HW, and Kim YHB (2017). Effect of Probiotic Feeding on Oxidative Stability and Meat Quality Attributes of Breast Muscle from Chickens Exposed to Chronic Heat Stress. *Meat and Muscle Biology*, 1 (3). [Article link](https://doi.org/10.22175/rmc2017.161) | DOI: <https://doi.org/10.22175/rmc2017.161>
- Chawke AP, Dakhore RS, and Shegokar SR (2021). Effect of Dietary Supplementation of Black Cumin (*Nigella sativa*) Seed Powder on Growth Performance and Economics of Giriraja Poultry. *International Journal of Current Microbiology and Applied Sciences*, 7(1): 56-63. [Article link](https://doi.org/10.20546/ijcmas.2021.1001.163) | DOI: <https://doi.org/10.20546/ijcmas.2021.1001.163>
- Chowdhury VS, and Furuse M (2013). Central regulation of physiological and metabolic responses in neonatal chicks. In *Chickens: Physiology, Diseases and Farming Practices*. Nova Science Publishers, Inc. pp. 105-124. [Article link](https://doi.org/10.1016/j.psj.2020.06.071) | DOI: <https://doi.org/10.1016/j.psj.2020.06.071>
- Cronje PB (2018). Essential role of methyl donors in animal productivity. *Animal Production Science*, 58 (4): 655-665. [Article link](https://doi.org/10.1071/AN15729) | DOI: <https://doi.org/10.1071/AN15729>
- Daghir N (2009). Nutritional strategies to reduce heat stress in broilers and broiler breeders. *Lohmann Information*, 44 (1), 6-15. [Article link](https://doi.org/10.1016/j.psj.2020.06.071) | DOI: <https://doi.org/10.1016/j.psj.2020.06.071>
- Daghir N.J (2008). Present status and future of the poultry industry in hot regions. *Poultry Production in Hot Climates: Second Edition*, CABI. Pp.1-11. [Article link](https://doi.org/10.1079/9781845932589.0001) | DOI: <https://doi.org/10.1079/9781845932589.0001>
- Doneria R, Dubey M, Gendley MK, Ramteke RC, and Pathak R (2020). Essential oils and Organic Acids: Feed Additive in Broiler Chickens. *International Journal of Current Microbiology and Applied Sciences*, 9(6): 3586-3593. [Article link](https://doi.org/10.20546/ijcmas.2020.906.422) | DOI: <https://doi.org/10.20546/ijcmas.2020.906.422>
- Dosoky WM, Zeweil HS, Ahmed MH, Zahran SM, Ali AM, Abdelsalam NR, and Naiel MAE (2021). The influences of Tylosine and licorice dietary supplementation in terms of the productive performance, serum parameters, egg yolk lipid profile, antioxidant and immunity status of laying Japanese quail under heat stress condition. *Journal of Thermal Biology*, 103015. [Article link](https://doi.org/10.1016/j.jtherbio.2021.103015) | DOI: <https://doi.org/10.1016/j.jtherbio.2021.103015>
- Tekce E and Gül M (2021). Effects of *Origanum syriacum* essential oil added in different levels to the diet of broilers under heat stress on performance and intestinal histology. *European Poultry Science*, 80: 1-11. [Article link](https://doi.org/10.1399/eps.2016.157) | DOI: <https://doi.org/10.1399/eps.2016.157>
- El-Kassas S, El-Naggar K, Abdo SE, Abdo W, Kirrella AAK, El-Mehaseeb I, and Abu El-Magd M (2019). Dietary supplementation with copper oxide nanoparticles ameliorates chronic heat stress in broiler chickens. *Animal Production Science*, 60 (2): 254-268. [Article link](https://doi.org/10.1071/AN18270) | DOI: <https://doi.org/10.1071/AN18270>
- El-Shoukary RDM, Darwish MHA, and Abdel-Rahman MAM (2014). Behavioral, Performance, Carcass Traits and Hormonal Changes of Heat Stressed Broilers Feeding Black and Coriander Seeds. *Advanced Veterinary Research*, 4(3): 97-101. [Article link](https://doi.org/10.1007/s11250-021-02554-0) | DOI: <https://doi.org/10.1007/s11250-021-02554-0>
- Elbaz AM, Ibrahim NS, Shehata AM, Mohamed NG, and Abdel-Moneim AME (2021). Impact of multi-strain probiotic, citric acid, garlic powder or their combinations on performance, ileal histomorphometry, microbial enumeration and humoral immunity of broiler chickens. *Tropical Animal Health and Production*, 53(1): 1-10. [Article link](https://doi.org/10.1007/s11250-021-02554-0) | DOI: <https://doi.org/10.1007/s11250-021-02554-0>
- Farfán C, Oliveros Y, and de Basilio V (2010). Effect of mineral supplying in water or feed on productive and physiological variables in broiler under heat stress conditions. *Zootecnia Tropical*, 28(3): 363-373. [Article link](https://doi.org/10.1007/s11250-021-02554-0) | DOI: <https://doi.org/10.1007/s11250-021-02554-0>
- Fathi MM, Ebeid TA, Al-Homidan I, Soliman NK, and Abou-Emera OK (2017). Influence of probiotic supplementation on immune response in broilers raised under hot climate. *British Poultry Science*, 58 (5): 512-516. [Article link](https://doi.org/10.1080/00071668.2017.1332405) | DOI: <https://doi.org/10.1080/00071668.2017.1332405>
- Gharaghani H, Shariatmadari F, and Torshizi MA (2015). Effect of fennel (*Foeniculum vulgare* mill.) used as a feed additive on the egg quality of laying hens under heat stress. *Brazilian Journal of Poultry Science*, 17: 199-207. [Article link](https://doi.org/10.1590/1516-635x1702199-208) | DOI: <https://doi.org/10.1590/1516-635x1702199-208>
- Ghasemi HA and Nari N (2020). Effect of supplementary betaine on growth performance, blood biochemical profile, and immune response in heat-stressed broilers fed different dietary protein levels. *Journal of Applied Poultry Research*, 29 (2): 301-313. [Article link](https://doi.org/10.1016/j.japr.2019.11.004) | DOI: <https://doi.org/10.1016/j.japr.2019.11.004>
- Gonçalves SA, Ferreira RA, Pereira IG, De Oliveira CC, Amaral PIS, Garbossa CAP, and Da Silva Fonseca L (2017). Behavioral and physiological responses of different genetic lines of free-range broiler raised on a semi-intensive system. *Journal of Animal Behaviour and Biometeorology*, 5 (4): 112-117. [Article link](https://doi.org/10.14269/2318-1265/jabb.v5n4p112-117) | DOI: <https://doi.org/10.14269/2318-1265/jabb.v5n4p112-117>
- Gorlov IF, Slozhenkina MI, Komarova ZB, Golovin VV, Krotova OE, Ivanov SM, and Rudkovskaya AV (2020). Mineral feed additive to prevent chickens' heat stress. *International Journal of Pharmaceutical Research*, 12 (3): 168-173. [Article link](https://doi.org/10.31838/ijpr/2020.12.03.030) | DOI: <https://doi.org/10.31838/ijpr/2020.12.03.030>

- Greene ES, Cauble R, Kadhim H, de Almeida Mallmann B, Gu I, Lee SO, and Dridi S (2021). Protective effects of the phyto-genic feed additive “comfort” on growth performance via modulation of hypothalamic feeding- and drinking-related neuropeptides in cyclic heat-stressed broilers. *Domestic Animal Endocrinology*, 74: 106487. [Article link | DOI: https://doi.org/10.1016/j.domaniend.2020.106487](https://doi.org/10.1016/j.domaniend.2020.106487)
- Hafeez A, Akram W, Sultan A, Konca Y, Ayasan T, Naz S, and Khan RU (2021). Effect of dietary inclusion of taurine on performance, carcass characteristics and muscle micro-measurements in broilers under cyclic heat stress. *Italian Journal of Animal Science*, 20(1): 872-877. [Article link | DOI: https://doi.org/10.1080/1828051X.2021.1921627](https://doi.org/10.1080/1828051X.2021.1921627)
- Hajati H, Hassanabadi A, Golian A, Nassiri-Moghaddam H, and Reza Nassiri M (2015). The effect of grape seed extract and vitamin C feed supplementation on some blood parameters and HSP70 gene expression of broiler chickens suffering from chronic heat stress. *Italian Journal of Animal Science*, 14 (3): 3273. [Article link | DOI: https://doi.org/10.4081/ijas.2015.3273](https://doi.org/10.4081/ijas.2015.3273)
- Hassan Mir S, Prakash Pal R, Mani V, Ahmed Malik T, Ojha L, and Yadav S (2018). Role of dietary minerals in heat-stressed poultry: A review. *Journal of Entomology and Zoology Studies*, 6, 820-826. [Article link | DOI: https://doi.org/10.1007/978-3-030-54462-1_7](https://doi.org/10.1007/978-3-030-54462-1_7)
- He S, Li S, Arowolo MA, Yu Q, Chen F, Hu R, and He J (2019). Effect of resveratrol on growth performance, rectal temperature and serum parameters of yellow-feather broilers under heat stress. *Animal Science Journal*, 90 (3): 401-411. [Article link | DOI: https://doi.org/10.1111/asj.13161](https://doi.org/10.1111/asj.13161)
- He W, Li P, and Wu G (2021). Amino Acid Nutrition and Metabolism in Chickens. *Advances in Experimental Medicine and Biology*, 1285: 109-131. [Article link | DOI: https://doi.org/10.1007/978-3-030-54462-1_7](https://doi.org/10.1007/978-3-030-54462-1_7)
- Holik V (2015). Management of laying hens under tropical conditions begins during the rearing period. *Lohmann Information*, 50(2):16-23. [Article link | DOI: https://doi.org/10.1007/978-3-030-54462-1_7](https://doi.org/10.1007/978-3-030-54462-1_7)
- Hosseini-Vashan SJ, and Piray AH (2021). Effect of dietary saffron (*Crocus sativus*) petal extract on growth performance, blood biochemical indices, antioxidant balance, and immune responses of broiler chickens reared under heat stress conditions. *Italian Journal of Animal Science*, 20(1): 1338-1347. [Article link | DOI: https://doi.org/10.1080/1828051X.2021.1921628](https://doi.org/10.1080/1828051X.2021.1921628)
- Humam AM, Loh TC, Foo HL, Izuddin WI, Zulkifli I, Samsudin AA, and Mustapha NM (2021). Supplementation of postbiotic RI11 improves antioxidant enzyme activity, upregulated gut barrier genes, and reduced cytokine, acute phase protein, and heat shock protein 70 gene expression levels in heat-stressed broilers. *Poultry Science*, 100(3): 100908. [Article link | DOI: https://doi.org/10.1016/j.psj.2020.12.011](https://doi.org/10.1016/j.psj.2020.12.011)
- Iyasere OS, Bateson M, Beard AP, and Guy JH (2021). Which factor is more important: Intensity or duration of episodic heat stress on broiler chickens? *Journal of Thermal Biology*, 102981. [Article link | DOI: https://doi.org/10.1016/j.jtherbio.2021.102981](https://doi.org/10.1016/j.jtherbio.2021.102981)
- Jimoh AA, Ahmed A, Agaie BM, Umaru MA, Ibrahim A, Jelani Ibitoye EB (2020). Ameliorative effects of *Citrullus lanatus* (watermelon) juice on survival, performance and lipid peroxidation status of heat-stressed broiler chickens. *Nigerian Journal of Animal Production*, 45 (4): 155-165. [Article link | DOI: https://doi.org/10.51791/njap.v45i4.551](https://doi.org/10.51791/njap.v45i4.551)
- Kang D and Shim K (2021). Early heat exposure effect on the heat shock proteins in broilers under acute heat stress. *Poultry Science*, 100 (3): 100964. [Article link | DOI: https://doi.org/10.1016/j.psj.2020.12.061](https://doi.org/10.1016/j.psj.2020.12.061)
- Khafar KR, Mojtahedin A, Rastegar N, Neytali MK, and Olfati A (2019). Dietary inclusion of thyme essential oil alleviative effects of heat stress on growth performance and immune system of broiler chicks. *Iranian Journal of Applied Animal Science*, 9 (3):509-517. [Article link | DOI: https://doi.org/10.1016/j.psj.2020.12.061](https://doi.org/10.1016/j.psj.2020.12.061)
- Khan RU, Naz S, and Dhama K (2014). Chromium: Pharmacological applications in heat-stressed poultry. *International Journal of Pharmacology*, 9 (3): 509-517. [Article link | DOI: https://doi.org/10.3923/ijp.2014.213.217](https://doi.org/10.3923/ijp.2014.213.217)
- Koh K and Macleod MG (1999). Effects of ambient temperature on heat increment of feeding and energy retention in growing broilers maintained at different food intakes. *British Poultry Science*, 40 (4): 511-516. [Article link | DOI: https://doi.org/10.1080/00071669987287](https://doi.org/10.1080/00071669987287)
- Kumbhar S, Khan AZ, Parveen F, Nizamani ZA, Siyal FA, El-Hack MEA, Huang K (2018). Impacts of selenium and vitamin E supplementation on mRNA of heat shock proteins, selenoproteins and antioxidants in broilers exposed to high temperature. *AMB Express*, 8 (1), 1-10. [Article link | DOI: https://doi.org/10.1186/s13568-018-0641-0](https://doi.org/10.1186/s13568-018-0641-0)
- Kuttappan V.A, Manangi M, Bekker M, Chen J, and Vazquez-Anon M (2021). Nutritional Intervention Strategies Using Dietary Antioxidants and Organic Trace Minerals to Reduce the Incidence of Wooden Breast and Other Carcass Quality Defects in Broiler Birds. *Frontiers in Physiology*, 12: 428. [Article link | DOI: https://doi.org/10.3389/fphys.2021.663409](https://doi.org/10.3389/fphys.2021.663409)
- Laġan C, Ribeiro AML, Kessler AM, Kratz LR, and Pinheiro CC (2007). Effects of the reduction of dietary heat increment on the performance, carcass yield, and diet digestibility of broilers submitted to heat stress. *Revista Brasileira de Cincia Avcola*, 9: 45-51. [Article link | DOI: https://doi.org/10.1590/S1516-635X2007000100007](https://doi.org/10.1590/S1516-635X2007000100007)
- Lartey KA, Kang DJ, Zhang QH, Shi CQ, Yang F, Lin HY, Chen JJ (2020). *Chromolaena odorata* as a potential feed additive bioresource to alleviate heat stress in chickens in the humid tropics. *World’s Poultry Science Journal*, 76 (1): 119-133. [Article link | DOI: https://doi.org/10.1080/00439339.2020.1729673](https://doi.org/10.1080/00439339.2020.1729673)
- Laudadio V, Dambrosio A, Normanno G, Khan R.U, Naz S, Rowghani E, and Tufarelli V (2012). Effect of reducing dietary protein level on performance responses and some microbiological aspects of broiler chickens under summer environmental conditions. *Avian Biology Research*, 5 (2), 88-92. [Article link | DOI: https://doi.org/10.3184/175815512X1335018071](https://doi.org/10.3184/175815512X1335018071)
- Liew PK, Zulkifli I, Hair-Bejo M, Omar AR, and Israfi DA (2003). Effects of early age feed restriction and heat conditioning on heat shock protein 70 expression, resistance to infectious bursal disease, and growth in male broiler chickens subjected to heat stress. *Poultry Science*, 82(12): 1879-1885. [Article link | DOI: https://doi.org/10.1093/ps/82.12.1879](https://doi.org/10.1093/ps/82.12.1879)
- Lin H, Jiao HC, Buyse J, and Decuyper E (2006). Strategies for preventing heat stress in poultry. *World’s Poultry Science Journal*, 62(1): 71-86. [Article link | DOI: https://doi.org/10.1079/WPS200585](https://doi.org/10.1079/WPS200585)
- Liu LL, He JH, Xie HB, Yang YS, Li JC, and Zou Y (2014). Resveratrol induces antioxidant and heat shock protein mRNA expression in response to heat stress in black-boned chickens. *Poultry Science*, 93 (1): 54-62. [Article link | DOI: https://doi.org/10.3382/ps.2013-03423](https://doi.org/10.3382/ps.2013-03423)
- Liu L, Ren M, Ren K, Jin Y, and Yan M (2020). Heat stress impacts on broiler performance: a systematic review and meta-analysis. *Poultry Science*, 99 (11): 6205-6211. [Article link | DOI: https://doi.org/10.1016/j.psj.2020.08.019](https://doi.org/10.1016/j.psj.2020.08.019)
- Liu M, Lu Y, Gao P, Xie X, Li D, Yu D, and Yu M (2020). Effect of curcumin on laying performance, egg quality, endocrine hormones, and immune activity in heat-stressed hens. *Poultry Science*, 99 (4): 2196-2202. [Article link | DOI: https://doi.org/10.1016/j.psj.2019.12.001](https://doi.org/10.1016/j.psj.2019.12.001)
- Liverpool-Tasie LSO, Sanou A, and Tambo JA (2019). Climate change adaptation among poultry farmers: evidence from Nigeria. *Climatic Change*, 157(3): 527-544. [Article link | DOI: https://doi.org/10.1007/s10584-019-02574-8](https://doi.org/10.1007/s10584-019-02574-8)

- Mack LA, Felver-Gant JN, Dennis RL, and Cheng HW (2013). Genetic variations alter production and behavioral responses following heat stress in 2 strains of laying hens, *Poultry Science*, 92(2): 285-294. [Article link](#) | DOI: <https://doi.org/10.3382/ps.2012-02589>
- Mahmoud UT, Abdel-Rahman MAM, Darwish MHA, Applegate TJ, and Cheng HW (2015). Behavioral changes and feathering score in heat stressed broiler chickens fed diets containing different levels of propolis. *Applied Animal Behaviour Science*, 166: 98-105. [Article link](#) | DOI: <https://doi.org/10.1016/j.applanim.2015.03.003>
- Mehaisen G.M.K, Ibrahim R.M, Desoky A.A, Safaa H.M, El-Sayed O.A, and Abass A.O (2017). The importance of propolis in alleviating the negative physiological effects of heat stress in quail chicks. *PLoS ONE*, 12 (10), e0186907. [Article link](#) | DOI: <https://doi.org/10.1371/journal.pone.0186907>
- Malekizadeh M, Moeini MM, Ghazi S (2013). The effects of different levels of ginger (*Zingiber officinale* Rosc) and turmeric (*Curcuma longa* Linn) rhizomes powder on some blood metabolites and production performance characteristics of laying hens. *Journal of agricultural science and technology*, 3(2): 301-305. [Article link](#) | DOI: <https://doi.org/10.1016/j.jst.2013.03.003>
- Min YN, Liu FX, Qi X, Ji S, Ma SX, Liu X, Gao YP (2018). Effects of methionine hydroxyl analog chelated zinc on laying performance, eggshell quality, eggshell mineral deposition, and activities of Zn-containing enzymes in aged laying hens. *Poultry Science*, 97 (10):3587-3593. [Article link](#) | DOI: <https://doi.org/10.3382/ps/pey203>
- Mohamed ASA, Lozovskiy AR, and Ali AMA (2019). Strategies to combat the deleterious impacts of heat stress through feed restrictions and dietary supplementation (vitamins, minerals) in broilers. *Journal of the Indonesian Tropical Animal Agriculture*, 44:155-166. [Article link](#) | DOI: <https://doi.org/10.14710/jitaa.44.2.155-166>
- Mujahid A (2011). Nutritional strategies to maintain efficiency and production of chickens under high environmental temperature. *Journal of Poultry Science*, 1105190115-1105190115. [Article link](#) | DOI: <https://doi.org/10.2141/jpsa.010115>
- Mushtaq T, Mirza MA, Athar M, Hooge DM, Ahmad T, Ahmad G Noreen U. (2007). Dietary sodium and chloride for twenty-nineto forty-two-day-old broiler chickens at constant electrolyte balance under subtropical summer conditions. *Journal of Applied Poultry Research*, 16 (2): 161-170. [Article link](#) | DOI: <https://doi.org/10.1093/japr.16.2.161>
- Mustafa MA and Wasman PH (2020). The impact of powders and oil additives of cinnamon and clove in quails diet as antistressor and antioxidant during hot months. *Iraqi Journal of Agricultural Sciences*, 51(3): 760-766. [Article link](#) | DOI: <https://doi.org/10.36103/ijas.v51i3.1031>
- Nawab A Li G, An L, wu J, Chao L, Xiao M, Ghani MW (2019). Effect of curcumin supplementation on TLR4 mediated non-specific immune responses in liver of laying hens under high-temperature conditions. *Journal of Thermal Biology*, 84: 384-397. [Article link](#) | DOI: <https://doi.org/10.1016/j.jtherbio.2019.07.003>
- Osti R, Bhattarai D, and Zhou D (2017). Climatic variation: Effects on stress levels, feed intake, and bodyweight of broilers. *Revista Brasileira de Ciencia Avicola*, 19: 489-496. [Article link](#) | DOI: <https://doi.org/10.1590/1806-9061-2017-0494>
- Petričević V, Lukić M, Škrbić Z, Rakonjac S, Dosković V, Petričević M, and Stanojković A (2018). The effect of using rosemary (*Rosmarinus officinalis*) in broiler nutrition on production parameters, slaughter characteristics, and gut microbiological population. *Turkish Journal of Veterinary and Animal Sciences*, 42(6): 658-664. [Article link](#) | DOI: <https://doi.org/10.3906/vet-1803-53>
- Prajapat UK, Jain D, Sharma T, Dhuria RK, Bothra T, Nehra R, Siyag SS (2020). Effect of supplementation of tulsi (*Ocimum sanctum*) and fenugreek (*trigonella foenum-graceum* L.) on liveability and economics of heat stressed broilers in arid zone of Rajasthan. *International Journal Of Livestock Research*, 8(10): 247-251. [Article link](#) | DOI: <http://doi.org/10.5455/ijlr.21080317050254>
- Raei A, Karimi A, and Sadeghi A (2020). Performance, antioxidant status, nutrient retention and serum profile responses of laying Japanese quails to increasing addition levels of dietary guanidinoacetic acid. *Italian Journal of Animal Science*, 42(6): 658-664. [Article link](#) | DOI: <https://doi.org/10.1080/1828051X.2019.1698325>
- Ranjan A, Sinha R, Devi I, Rahim A, and Tiwari S (2019). Effect of Heat Stress on Poultry Production and their Managemental Approaches. *International Journal of Current Microbiology and Applied Sciences*, 8 (02): 1548-1555. [Article link](#) | DOI: <https://doi.org/10.20546/ijcmas.2019.802.181>
- Rao H and Gurrām S (2021). Effect of herbal feed additives on performance, immunity, serum parameters and E. coli counts of broilers under heat stress. *Indian Journal of Animal Nutrition*, 38 (1): [Article link](#) | DOI: <https://doi.org/10.5958/2231-6744.2021.00009.8>
- Rao SVR, Prakash B, Raju MVLN, Panda AK, Kumari RK, and Reddy EPK (2016). Effect of Supplementing Organic Forms of Zinc, Selenium and Chromium on Performance, Anti-Oxidant and Immune Responses in Broiler Chicken Reared in Tropical Summer. *Biological Trace Element Research*, 172 (2): 511-520. [Article link](#) | DOI: <https://doi.org/10.1007/s12011-015-0587-x>
- Ratriyanto A, Indreswari R, and Nuhriawangsa AMP (2017). Effects of dietary protein level and betaine supplementation on nutrient digestibility and performance of Japanese quails. *Brazilian journal of poultry science*, 19: 445-454. [Article link](#) | DOI: <https://doi.org/10.1590/1806-9061-2016-0442>
- Ratriyanto A, Indreswari R, and Sunarto (2014). Effects of protein levels and supplementation of methyl group donor on nutrient digestibility and performance of broiler chickens in the tropics. *International Journal of Poultry Science*, 13 (10): 575-581. [Article link](#) | DOI: <https://doi.org/10.3923/ijps.2014.575.581>
- Ratriyanto A and Mosenthin R (2018). Osmoregulatory function of betaine in alleviating heat stress in poultry. *Journal of Animal Physiology and Animal Nutrition*, 102 (6): 1634-1650. [Article link](#) | DOI: <https://doi.org/10.1111/jpn.12990>
- Ryder AA, Feddes JJR, and Zuidhof MJ (2004). Field study to relate heat stress index to broiler performance. *Journal of Applied Poultry Research*, 13(3): 493-499. [Article link](#) | DOI: <https://doi.org/10.1093/japr/13.3.493>
- Sadeghi AA and Moghaddam M (2018). The effects of turmeric, cinnamon, ginger and garlic powder nutrition on antioxidant enzymes' status and hormones involved in energy metabolism of broilers during heat stress. *Iranian Journal of Applied Animal Science*, 8 (1): 125-130. [Article link](#) | DOI: <https://doi.org/10.1016/j.ijas.2018.03.003>
- Sahin K, Orhan C, Tuzcu M, Ali S, Sahin N, and Hayirli A (2010). Epigallocatechin-3-gallate prevents lipid peroxidation and enhances antioxidant defense system via modulating hepatic nuclear transcription factors in heat-stressed quails. *Poultry Science*, 89 (10): 2251-2258. [Article link](#) | DOI: <https://doi.org/10.3382/ps.2010-00749>
- Sahin, Kazim, Sahin N, Onderci M, Yaralioglu S, and Kucuk O (2001). Protective role of supplemental vitamin E on lipid peroxidation, vitamins E, A and some mineral concentrations of broilers reared under heat stress. *Veterinary Medicine*, 46 (5): 140-144. [Article link](#) | DOI: <https://doi.org/10.17221/7870-VETMED>
- Saiz del Barrio A, Mansilla WD, Navarro-Villa A, Mica JH, Smeets JH, den Hartog LA, and García-Ruiz AI (2020). Effect of mineral and vitamin C mix on growth performance and blood corticosterone concentrations in heat-stressed broilers. *Journal of Applied Poultry Research*, 29 (1): 23-33. [Article link](#) | DOI: <https://doi.org/10.1016/j.japr.2019.11.001>
- Saleh AA, Kirrella AA, Dawood MAO, and Ebeid TA (2019). Effect of dietary inclusion of cumin seed oil on the performance, egg quality, immune response and ovarian development in laying hens under high ambient temperature. *Journal of Animal Physiology and Animal Nutrition*, 103 (6): 1810-1817. [Article link](#) | DOI: <https://doi.org/10.1111/jpn.13206>

- Sandner G, Mueller AS, Zhou X, Stadlbauer V, Schwarzingler B, Schwarzingler C, Weghuber J (2020). Ginseng extract ameliorates the negative physiological effects of heat stress by supporting heat shock response and improving intestinal barrier integrity: Evidence from studies with heat-stressed Caco-2 cells, C. Elegans and Growing Broilers. *Molecules*, 25(4): 835. [Article link](https://doi.org/10.3390/molecules25040835) | DOI: <https://doi.org/10.3390/molecules25040835>
- Santos Dalólio F, Teixeira Albino LF, Nogueira Da Silva J, Kyonara Alves Tenório Fireman A, Burin ÁM, Busanello M, and Ribeiro V (2021). Dietary chromium-methionine supplementation and broiler (22-43 days) responses during heat stress. 1. Growth performance and carcass yield, metabolisable energy and serum biochemistry. *Animal Production Science*, 61 (6): 586-595. [Article link](https://doi.org/10.1071/AN20140) | DOI: <https://doi.org/10.1071/AN20140>
- Santos MM, Souza-Junior JBF, Queiroz JPAF, Costa MKO, Lima HFF, Arruda AMV, and Costa LLM (2019). Broilers' behavioural adjustments when submitted to natural heat stress and fed different maize particle sizes in the diet. *Journal of Agricultural Science*, 157(9-10): 743-748. [Article link](https://doi.org/10.1017/S0021859620000131) | DOI: <https://doi.org/10.1017/S0021859620000131>
- Saracila M, Panaite T.D, Papuc C.P, and Criste R.D (2021). Heat stress in broiler chickens and the effect of dietary polyphenols, with special reference to Willow (*Salix* spp.) bark supplements—A review. *Antioxidants*, 10(5): 686. [Article link](https://doi.org/10.3390/antiox10050686) | DOI: <https://doi.org/10.3390/antiox10050686>
- Sarwar M and Ahmad T (2007). Dietary electrolyte balance: implications in heat stressed broilers. *World's Poultry Science Journal*, 62 (4): 638-653. [Article link](https://doi.org/10.1017/wps2006118) | DOI: <https://doi.org/10.1017/wps2006118>
- Shahzad K, Sultan M, Bilal M, Ashraf H, Farooq M, Miyazaki T, Hussain MI (2021). Experiments on energy-efficient evaporative cooling systems for poultry farm application in Multan (Pakistan). *Sustainability (Switzerland)*, 13(5): 2836. [Article link](https://doi.org/10.3390/su13052836) | DOI: <https://doi.org/10.3390/su13052836>
- Siddiqui S.H, Kang D, Park J, Khan M, Belal S.A, Shin D, Shim K (2021). Altered relationship between gluconeogenesis and immunity in broilers exposed to heat stress for different durations. *Poultry Science*, 100(18): 101274. [Article link](https://doi.org/10.1016/j.psj.2021.101274) | DOI: <https://doi.org/10.1016/j.psj.2021.101274>
- Slawinska A, Zampiga M, Sirri F, Meluzzi A, Bertocchi M, Tavaniello S and Maiorano G (2020). Impact of galactooligosaccharides delivered in ovo on mitigating negative effects of heat stress on performance and welfare of broilers. *Poultry Science*, 99 (1): 407-415. [Article link](https://doi.org/10.3382/ps/pez512) | DOI: <https://doi.org/10.3382/ps/pez512>
- Sohail MU, Hume ME, Byrd JA, Nisbet DJ, Ijaz A, Sohail A, Shabbir MZ, Rehman H (2012). Effect of supplementation of prebiotic mannan-oligosaccharides and probiotic mixture on growth performance of broilers subjected to chronic heat stress. *Poultry science*, 91(9):2235-2240. [Article link](https://doi.org/10.3382/ps.2012-02182) | DOI: <https://doi.org/10.3382/ps.2012-02182>
- Song DJ and King AJ (2015). Effects of heat stress on broiler meat quality. *World's Poultry Science Journal*, 71(4):701-709. [Article link](https://doi.org/10.1017/S0043933915002421) | DOI: <https://doi.org/10.1017/S0043933915002421>
- Srinivasa Rao G, Gopinath KA, Prasad JVNS, Prasannakumar, and Singh AK (2016). Climate Resilient Villages for Sustainable Food Security in Tropical India: Concept, Process, Technologies, Institutions, and Impacts. *Advances in Agronomy*, 140: 101-214. [Article link](https://doi.org/10.1016/bs.agron.2016.06.003) | DOI: <https://doi.org/10.1016/bs.agron.2016.06.003>
- Surai PF, Kochish II, Romanov MN, and Griffin DK (2019). Nutritional modulation of the antioxidant capacities in poultry: the case of vitamin E. *Poultry Science*, 98 (9): 4030-4041. [Article link](https://doi.org/10.3382/ps/pez072) | DOI: <https://doi.org/10.3382/ps/pez072>
- Tamiru M, Ashagrie S, Alkhtib A, Getachew M, Demeke S, Hassen W, Burton E (2020). Performance of broilers and layers supplemented with *Moringa stenopetala* leaf meal under hot humid tropical conditions. *Animal Production Science*, 60 (17): 1987-1994. [Article link](https://doi.org/10.1071/AN19671) | DOI: <https://doi.org/10.1071/AN19671>
- Tang S, Yin B, Xu J, and Bao E (2018). Rosemary reduces heat stress by inducing CRYAB and HSP70 expression in broiler chickens. *Oxidative Medicine and Cellular Longevity*, 2018: Article ID 7014126. [Article link](https://doi.org/10.1155/2018/7014126) | DOI: <https://doi.org/10.1155/2018/7014126>
- Tolba SA, Magnuson AD, Sun T, and Lei XG (2020). Dietary supplemental microalgal astaxanthin modulates molecular profiles of stress, inflammation, and lipid metabolism in broiler chickens and laying hens under high ambient temperatures. *Poultry Science*, 99 (10): 4853-4860. [Article link](https://doi.org/10.1016/j.psj.2020.05.022) | DOI: <https://doi.org/10.1016/j.psj.2020.05.022>
- Wan X, Ahmad H, Zhang L, Wang Z, and Wang T (2018). Dietary enzymatically treated *Artemisia annua* L. improves meat quality, antioxidant capacity and energy status of breast muscle in heat-stressed broilers. *Journal of the Science of Food and Agriculture*, 98 (10): 3715-3721. [Article link](https://doi.org/10.1002/jsfa.8879) | DOI: <https://doi.org/10.1002/jsfa.8879>
- Wankhade PR, Talokar AJ, Patel B, and Verma D (2020). Management Interventions to Reduce Heat Stress in Chickens. *Quarterly Research Journal of Plant and Animal Sciences*, 35: 110-113. [Article link](https://doi.org/10.18805/bkap203) | DOI: <https://doi.org/10.18805/bkap203>
- Wasti S, Sah N, Lee CN, Jha R, and Mishra B (2021). Dietary supplementation of alpha-lipoic acid mitigates the negative effects of heat stress in broilers. *PLoS ONE*, 16 (7): e0254936. [Article link](https://doi.org/10.1371/journal.pone.0254936) | DOI: <https://doi.org/10.1371/journal.pone.0254936>
- Wasti S, Sah N, Singh A.K, Lee C.N, Jha R, and Mishra B (2021). Dietary supplementation of dried plum: a novel strategy to mitigate heat stress in broiler chickens. *Journal of Animal Science and Biotechnology*, 12 (1): 1-17. [Article link](https://doi.org/10.1186/s40104-021-00571-5) | DOI: <https://doi.org/10.1186/s40104-021-00571-5>
- Wu QJ, Liu N, Wu XH, Wang GY, and Lin L (2018). Glutamine alleviates heat stress-induced impairment of intestinal morphology, intestinal inflammatory response, and barrier integrity in broilers. *Poultry Science*, 97 (8): 2675-2683. [Article link](https://doi.org/10.3382/ps/pey123) | DOI: <https://doi.org/10.3382/ps/pey123>
- Xue B, Song J, Liu L, Luo J, Tian G, and Yang Y (2017). Effect of epigallocatechin gallate on growth performance and antioxidant capacity in heat-stressed broilers. *Archives of Animal Nutrition*, 71 (5): 362-372. [Article link](https://doi.org/10.1080/1745039X.2017.1355129) | DOI: <https://doi.org/10.1080/1745039X.2017.1355129>
- Yang C, Luo P, Chen S, jian, Deng Z, chao, Fu X, liang, Xu, D. ning, Liu, W. jun (2021). Resveratrol sustains intestinal barrier integrity, improves antioxidant capacity, and alleviates inflammation in the jejunum of ducks exposed to acute heat stress. *Poultry Science*, 100(11): 101459. [Article link](https://doi.org/10.1016/j.psj.2021.101459) | DOI: <https://doi.org/10.1016/j.psj.2021.101459>
- Zhang J, Bai K.W, He J, Niu Y, Lu Y, Zhang L, and Wang T (2018). Curcumin attenuates hepatic mitochondrial dysfunction through the maintenance of thiol pool, inhibition of mtDNA damage, and stimulation of the mitochondrial thioredoxin system in heat-stressed broilers. *Journal of Animal Science*, 96 (3): 867-879. [Article link](https://doi.org/10.1093/jas/sky009) | DOI: <https://doi.org/10.1093/jas/sky009>
- Zhang P, Yan T, Wang X, Kuang S, Xiao Y, Lu W, and Bi D (2017). Probiotic mixture ameliorates heat stress of laying hens by enhancing intestinal barrier function and improving gut microbiota. *Italian Journal of Animal Science*, 16(2): 292-300. [Article link](https://doi.org/10.1080/1828051X.2016.1264261) | DOI: <https://doi.org/10.1080/1828051X.2016.1264261>
- Zhao GP, Han MJ, Zheng MQ, Zhao JP, Chen JL, and Wen J (2011). Effects of dietary vitamin E on immunological stress of layers and their offspring. *Journal of Animal Physiology and Animal Nutrition*, 95 (3): 343-350. [Article link](https://doi.org/10.1111/j.1439-0396.2010.01060.x) | DOI: <https://doi.org/10.1111/j.1439-0396.2010.01060.x>