

DOI: https://dx.doi.org/10.51227/ojafr.2025.35

# EFFECT OF DIETARY SUPPLEMENTATION WITH HYDROPONIC GREEN BARLEY FODDER ON RABBIT PERFORMANCE, ANTIOXIDANT CAPACITY, AND TOTAL PHENOLIC COMPOUNDS IN MEAT

Emilio A. DELIS-HECHAVARRIA<sup>1,2</sup>, Ramon G. GUEVARA-GONZALEZ<sup>3</sup>, José G. GOMEZ-SOTO<sup>4</sup>, Rosalía V. OCAMPO-VELAZQUEZ<sup>3</sup>, Irineo TORRES-PACHECO<sup>3</sup>, and Ixchel PAROLA-CONTRERAS<sup>2,5</sup>

ABSTRACT: Rabbit meat stands out as a nutritious low-fat protein source rich in essential amino acids. vitamins, and minerals, yet its production faces challenges from high feed costs and environmental impacts. Hydroponic green barley fodder (HGBF) emerges as a promising sustainable alternative, offering rapid growth, enhanced nutrient profiles including antioxidants, and reduced resource demands. This study aimed to evaluate the effects of dietary HGBF supplementation, with or without hydrogen peroxide (H2O2) seed pretreatment, on rabbit performance, meat antioxidant capacity, and total phenolic compounds. Sixty-three postweaned New Zealand rabbits (initial weight: 606 g) were divided into three groups for 42 days: T1 (100% pellets, control), T2 (50% pellets + 50% HGBF at 75 mM H<sub>2</sub>O<sub>2</sub> pre-treatment), and T3 (50% pellets + 50% HGBF), All treatments were ad libitum. Carcass yields were assessed post-slaughter, while meat samples underwent 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS\*+) assays for antioxidant activity, enzymatic evaluations (catalase, superoxide dismutase, phenylalanine ammonia-lyase), and Folin-Ciocalteu method for phenolics. Results revealed no significant differences in hot or cold carcass weights across treatments, averaging below export standards due to the rabbits' young age. However, T2 diet significantly boosted total phenolic content (P < 0.05) and antioxidant capacity via DPPH (97.52% inhibition) and ABTS<sup>++</sup> (84.07% inhibition) compared to T1 and T3 groups with lower catalase activity indicating reduced oxidative stress. Enzymatic activities for superoxide dismutase and phenylalanine ammonia-lyase remained unaffected. These findings underscore HGBF with H<sub>2</sub>O<sub>2</sub> pretreatment as an effective strategy to enhance meat quality through elevated phenolics and antioxidants, fostering oxidative stability and nutritional value while promoting eco-friendly rabbit farming practices for broader livestock sustainability.

**Keywords:** Antioxidants, Carcass yield, Hydroponic green barley fodder, Oxidative stability, Rabbit meat, Total phenolic compounds.

# INTRODUCTION

Rabbit meat is consumed all over the world, being delicious, appetizing and high quality. It is composition is mainly lipids, proteins and water, with high levels of micronutrients, essential amino acids, vitamins and minerals. Additionally, it has low uric acid, purines, and fat (Dalle Zotte and Szendro, 2011). However, it has been studied that the composition of the lipids depends on the type of rabbits feeding, being reflected in carcass composition (Oo et al., 2018). Furthermore, rabbit production is economically relevant for their sporting utility and the manufacture of skins that are used in multiple products (Pelletier et al., 2019). Furthermore, this species, the New Zealand rabbit, is commonly utilized in biological experiments within educational institutions and hospitals, serving as a valuable model for nutritional and antioxidant research (Song et al., 2020). Rising rabbits demand large volume of raw plant material to fattening (Wang et al., 2020). Fattening rabbits requires substantial plant-based feed, including tubers, cereals, fruits, and vegetables. Wild rabbits mainly feed of herbaceous and grasses, which contain considerable quantities of cellulose (Grioui et al., 2021).

In 2015, global cereal production reached 10 million tons; about 60% of world consumption of coarse grains (barley, maize, oats, quinoa, rye, sorghum, tef, and millet) used for livestock feed (FAO, 2021). Cereals also are produced for fodder livestock, since they can be converted directly into protein usable by animals, lowering the cost of feed, accelerating around 25% of growing and diminish time of fattening (Miah et al., 2020). Barley, constituting 65% of



PII: S222877012500035-15
Received: June 28, 2025
Revised: October 30, 2025
Accepted: November 02, 2025

<sup>&</sup>lt;sup>1</sup>CONAHCYT-Estancias posdoctorales, Universidad Nacional Autónoma de México, Universidad 3000, CDMX, 04510, Mexico

<sup>&</sup>lt;sup>2</sup>Red de Investigación Multidisciplinaria Integral UAEM 6304/REPD2020, El Tejocote, Texcoco, 56259, Mexico

<sup>&</sup>lt;sup>3</sup>Centro de Investigación Aplicada en Biosistemas (CIAB), Facultad de Ingeniería-Campus Amazcala, Universidad Autónoma de Querétaro, Carretera Chichimequillas S/N, El Marqués, 76265, Mexico

<sup>&</sup>lt;sup>4</sup>Facultad de Ciencias Naturales, Universidad Autónoma de Querétaro, A. de las Ciencias S/N, Querétaro, 76230, Mexico

<sup>&</sup>lt;sup>5</sup>TecNM/Tecnológico de Estudios Superiores de Chimalhuacán, División de Ingeniería Industrial, Ingeniería de Procesos Sustentables ITESCHIM-CA-02, Primavera S/N, Chimalhuacán, 55330, Mexico

Email: ixchelparola@teschi.edu.mx

Supporting Information

livestock feed. Whole grain barley includes flavonoids, folates, lignans, phenolic acids, phytosterols and tocoles, with the predominant vitamin E (tocol), flavonoids, phenols, sterols and vitamins. These phytochemicals offer strong antioxidant capacity, useful for reducing free radicals. In addition, this specie is rich in dietary fiber made up of soluble and insoluble fibers (Abouelezz et al., 2019). Hydroponic Green Barley Fodder (HGBF) yields 7–10 kg per kg of grain in 8–15 days, with increased crude protein, and calcium content; additionally, acid detergent fiber, neutral detergent fiber, calcium content and crude protein are increased (Singh et al., 2019).

Furthermore, rabbit meat production requires diets that mixture between 100 to 200 g of HGBF and a daily average of 50 g of concentrates, per animal, for optimal production (Miah et al., 2020). Furthermore, HGBF includes great fiber content, that it is necessary for rabbits' digestion and nutrition (Delis-Hechavarria et al., 2021). Besides, Selim et al. (2021) obtained a final live weight gain (1,533 g) with a supplementation of 15 g/kg with *Moringa oleifera*. Additionally, Abouelezz et al. (2019) had the highest body weight in male rabbits using concentrated plus HGBF and the feed intake was similar in all treatments. Indeed, the mixture of two feeds allowed the absorption of nutrients for growth, which could have a benign effect on the characteristic of rabbit meat (Dalle Zotte et al., 2018). Meat is susceptible of losing its nutritional properties due to the action of the oxidation process on lipids to rather than the degradation process by bacteria (Chen et al., 2017). This deterioration is observed in the discoloration of the meat. For example, Beefs are most susceptible to this oxidation process, reducing quality and shelf life (Bouzaida et al., 2021). The chemical processes involved in this process are the water retention capacity, rancid odors due to the formation of aldehydes and ketones, reduced solubility of proteins (Kim et al., 2019). However, studies on the concentrated diet plus HGBF are very scarce. Therefore, in this manuscript the effect of two combinations of barley fodder one with seeds pre germination on 75 mM hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and barley fodder with water supplemented plus 50% pellet, were evaluated on yield carcass and rabbit meat antioxidant capacity.

## **MATERIALS AND METHODS**

# Plant material and growth conditions

Hydroponic Green Barley Fodder (HGBF) was cultivated at Innovation and development Green (IDGREEN) farm, La Norita, Queretaro, México (20°32'58.6"N -100°30'10.6"W; altitude 1823 m) in January-February 2021. Barley seeds (Hordeum vulgare L var. Cantabria) were soaked in water and 75 mM hydrogen peroxide concentration for 24 h. It drained and placed on trays of 54 × 35 × 7 cm. The trays were stored in a chamber fodder for 11 days at 25°C, 14 h light:10 h dark photoperiod, and relative humidity of 80%. Fodder was watered only with water, six times daily for two min each one. Rabbits were fed with HGBF (leaves, roots and seeds) immediately after harvesting, including (Morales et al., 2010).

# **Animal and materials**

Sixty-three post-weaned New Zealand rabbits (average initial body weight (BW):  $0.606 \pm 0.01$  kg) were housed in four wire cages ( $76 \times 48 \times 30$  cm) containing three rabbits for each treatment. The temperature in the rabbitry was  $15 \pm 2$  °C,  $41.10 \pm 10.68\%$  relative air humidity and 12-h light: 12-h dark photoperiod. Rabbits had free access to clean water. The 42 days study included three fed diets: T1 (100 % pellets), T2 (50% pellets + HGBF at 75mM H $_2O_2$ ), T3 (50% pellets + HGBF ad libitum). All treatments were ad libitum. Rabbits were weighed and tranquillized with azaperone (2 mg/kg BW), anaesthetized with isoflurane (inhaled) and euthanized using a cardiac overdose of sodium pentobarbital (50 mg/kg BW) (Gómez Soto et al., 2018).

# **Ethical regulations**

The study was performed at DGREEN farm. The protocol was reviewed and approved by the Autonomous University of Queretaro's Bioethics Committee (authorization: 90-FI-2017) of the Engineering Science Faculty's. The handling of the experimental animals was carried out conforming to the Mexican Official Norm 'NOM-062-Z00-1999' guidelines (Norma Oficial Mexicana, 1999) and the International Guiding Principles for Biomedical Research Involving Animals (CIOMS/ICLAS 2012).

# Sampling

Analyses were carried out at the Animal Nutrition Laboratory, Faculty of Natural Sciences, Autonomous University of Queretaro, Queretaro, Mexico. The stomach, jejunum, ileum, cecum, and colon were dissected, and emptied contents measured a potentiometer (Model PH211, Hanna Instruments®). Jejunum-ileum, liver, gallbladder, stomach, colon-rectum and cecum sections were collected, washed with saline solution, weighed, and preserved in 10% neutral formalin solution for further analyses (Gómez Soto et al., 2018).

# Antioxidant activity (AOx)

Antioxidant properties of rabbit meat from each treatment were assessed using the 1,1-diphenyl-2-picrylhydrazyl

(DPPH •) and 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS •+) assays.

# DPPH • Radical Scavenging Activity (DRSA) Assay

The DPPH• assay was adapted from Chen et al. (2012) with modifications. A 0.5 mL sample solution's aliquot (600  $\mu$ g/mL) was combined with 0.5 mL of 0.1 mM DPPH• in methanol. The mixture was vigorously mixed, incubated in darkness at ambient temperature for 30 min, and absorbance was recorded at 515 nm. Methanol replaced DPPH• for the blank, and deionized water replaced the sample for the control. The DRSA percentage was calculated using the equation (1): DRSA(%) = [1-(A\_{sample}-A\_{blank}/A\_{control})] x 100 (1)

where A<sub>sample</sub> is the sample absorbance, A<sub>blank</sub> is the blank absorbance, and A<sub>control</sub> is the control absorbance.

# ABTS\*\* radical scavenging activity (ARSA) assay

The ARSA assay was conducted following Re et al. (1999) with adjustments. ABTS<sup>\*+</sup> radicals were produced by reacting 7 mM ABTS<sup>\*+</sup> with 2.45 mM potassium persulfate, incubating in darkness for 12 h at room temperature. The ABTS<sup>\*+</sup> solution was diluted in 0.15 M phosphate-buffered saline (PBS, pH 7.4) to achieve an absorbance of ~0.7 at 734 nm. Then, 3 mL of diluted ABTS<sup>\*+</sup> was mixed with 150  $\mu$ L of sample (600  $\mu$ g/mL), and absorbance was measured under dark conditions. PBS replaced ABTS<sup>\*+</sup> for the blank, and deionized water served as the control. ARSA was calculated as (2): ARSA(%) = [1-(A<sub>sample</sub>-A<sub>blank</sub>/A<sub>control</sub>)]x100 (2)

where A<sub>sample</sub> is the sample absorbance, A<sub>blank</sub> is the blank absorbance, and A<sub>control</sub> is the control absorbance.

# **Enzyme activity assays**

Enzyme activities in rabbit meat were evaluated to assess oxidative stress responses, focusing on catalase (CAT), superoxide dismutase (SOD), and phenylalanine ammonia-lyase (PAL).

## Samples preparation of enzymatic assays

Lyophilized rabbit meat samples (0.5 g) were homogenized in 1 mL of chilled extraction buffer and centrifuged at 13,000 rpm for 20 min at 4°C. The resulting supernatant was used for enzyme activity measurements. Protein content was quantified using the Bradford (1976) method, with bovine serum albumin (Sigma-Aldrich, St. Louis, MO, USA) as the standard.

# Catalase activity assay

Catalase (CAT; EC 1.11.1.6) activity was measured spectrophotometrically by monitoring the rate of  $H_2O_2$  decomposition at 240 nm, following Afiyanti and Chen (2014). The reaction mixture contained 0.95 mL of 50 mM potassium phosphate buffer (pH 8.0), 0.05 mL of enzymatic extract, and 0.1 mL of 100 mM  $H_2O_2$ . Absorbance changes were recorded for 1 min at 25 °C. One CAT unit decomposes 1  $\mu$ mol of  $H_2O_2$  per minute at pH 8.0. Results were expressed as U/mg protein.

# Superoxide dismutase activity assay

Superoxide dismutase (SOD; EC 1.15.1.1) activity was determined by measuring the inhibition of nitro blue tetrazolium (NBT) photochemical reduction, as described by Hayat et al. (2018). The reaction mixture included 1.5 mL of 50 mM potassium phosphate buffer (pH 7.8), 0.3 mL of 0.1 mM EDTA, 0.3 mL of 0.13 M methionine, 0.3 mL of 0.02 mM riboflavin, 0.05 mL of enzymatic extract, and 0.25 mL of distilled water. The mixture was exposed to fluorescent light (86.86 µmol/m²-s) for 20 min, and absorbance was measured at 560 nm. One SOD unit inhibits 50% of NBT reduction at pH 7.8 and 25°C. Results were reported as U/mg protein.

# Phenylalanine ammonia-lyase (PAL) activity

PAL activity (EC 4.3.1.5) was assessed using the method of Toscano et al. (2018). The assay measured cinnamic acid production from L-phenylalanine at 290 nm. The reaction mixture consisted of 2.3 mL of 0.1 M borate buffer (pH 8.8) with 10 mM L-phenylalanine and 0.2 mL of enzymatic extract. After incubation at  $40^{\circ}$ C for 1 h, the reaction was stopped with 0.5 mL of 5 N HCl. Absorbance was recorded at 290 nm. One PAL unit releases 1 µmol of cinnamic acid per minute at pH 8.8 and  $40^{\circ}$ C. Results were expressed as U/mg protein.

# **Total polyphenol content (TPC)**

Lyophilized rabbit meat samples (1 g) were pulverized in liquid nitrogen and extracted with a methanol:water mixture (1:10 w/v). The suspension was sonicated (Bransonic M2800-CPX-HE, Emerson, Ferguson, MO, USA) for 30 min, then centrifuged at 10,000 rpm for 15 min at 4°C. Supernatants were stored at 4°C in dark vials. Total phenols were quantified using the Folin-Ciocalteu method (Vergara-Castañeda et al., 2010) with gallic acid as the standard. A 140  $\mu$ L extract was mixed with 460  $\mu$ L of distilled water and 250  $\mu$ L of 1 N Folin-Ciocalteu reagent. After 5 min, 1250  $\mu$ L of 20% (w/v) sodium carbonate was added. The mixture was vortexed, incubated in darkness for 2 h, and absorbance was measured at 760 nm. Results were expressed as mg gallic acid equivalents per gram of dry matter (mg GAE/g DM).

## Statistical analysis

Rabbits were randomly divided into three dietary treatment groups, 21 animals being assigned to each one of the treatments (T1, T2, and T3, described in Animal and materials section) and each group having three replications with seven rabbits in each replication. Data were analyzed as a randomized complete block design. Data were subjected to analysis of variance (ANOVA) and Tukey tests ( $\alpha$  < 0.05) were using the Statgraphics® Centurion XVI statistical software (StatPoint Technologies Inc., Bedford, MA, USA, 2010).

## **RESULTS AND DISCUSSION**

#### Productive behavior of rabbit meat

The study determinates the productive performance of rabbit meat across three dietary treatments—100% pellets, 50% pellets + 50% HGBF at 75 mM  $H_2O_2$ , and 50% pellets + 50% HGBF—over a 42-day period, with results presented in Table 1. Live weight no was significantly different compared to the HGBF-supplemented groups, indicating a potential influence of pellet-based diets on growth. Stomach pH and organ weight showed no significant differences across treatments. Jejunum pH and length exhibited slight variations, though not statistically significant (P > 0.05). Cecum pH and length, colon-rectum pH, length, and liver relative weight remained consistent across groups, with no significant treatment effects (P > 0.05), suggesting that HGBF supplementation maintains gastrointestinal and organometric stability.

Parameters	Pellets 100%	Pellets 50% + 50% HGBF at 75 mM H₂O₂	Pellets 50% + 50% HGBF	
Alive weight (kg)	1.71 ± 0.09a	1.63 ± 0.16a	1.66 ± 0.15 <sup>a</sup>	
Stomach pH	1.59 ± 0.34 <sup>a</sup>	2.07 ± 1.22 <sup>a</sup>	1.91 ± 0.37ª	
Stomach organ empty weight (g)	19.58 ± 1.87a	20.45 ± 0.82°	20.85 ± 1.82ª	
Jejunum-ileum pH	7.33 ± 0.21 <sup>a</sup>	7.91 ± 1.07ª	7.43 ± 0.36 <sup>a</sup>	
Jejunum-ileum organ empty weight (g)	34.50 ± 5.66 <sup>a</sup>	34.73 ± 1.66a	38.27 ± 2.17a	
Jejunum-ileum length (cm)	205.50 ± 46.75 <sup>a</sup>	225.50 ± 16.05 <sup>a</sup>	207.00 ± 49.28a	
Cecum pH	5.85 ± 0.08 <sup>a</sup>	6.02 ± 0.18 <sup>a</sup>	5.96 ± 0.14 <sup>a</sup>	
Cecum organ empty weight (g)	24.60 ± 4.47a	21.30 ± 2.85a	25.13 ± 4.02a	
Cecum length (cm)	41.25 ± 3.03 <sup>a</sup>	40.25 ± 2.86a	39.25 ± 0.83°	
Colon-rectum pH	6.41 ± 0.27a	6.57 ± 0.14 <sup>a</sup>	6.46 ± 0.24 <sup>a</sup>	
Colon-rectum organ empty weight (g)	28.93 ± 5.36 <sup>a</sup>	27.85 ± 1.12a	28.95 ± 2.49a	
Colon-rectum length (cm)	95.25 ± 3.34a	95.75 ± 6.26 <sup>a</sup>	96.50 ± 4.56a	
Liver relative weight (g)	63.00 ± 11.42°	64.15 ± 11.23 <sup>a</sup>	62.83 ± 14.48a	
Gallbladder weight (g)	0.53 ± 0.13a	0.70 ± 0.27a	0.65 ± 0.32a	


In recent years, interest in the consumption of nutraceutical products has increased in the market (Cesari et al., 2018). Meat is considered a functional food with many bioactive compounds; specifically, rabbit meat has important nutritional and dietary properties to improve and support health (Dal Bosco et al., 2018). Figure 1 illustrates the hot and cold carcass weights and yields for the three treatments.

No significant differences were observed in hot or cold carcass weights across the treatments (T1, T2, T3). Average carcass weights were below typical export standards, attributed to the rabbits' young age at slaughter (initial age: 35 days; final age: ~77 days). This suggests that partial substitution of pellets with HGBF (with or without H<sub>2</sub>O<sub>2</sub> pre-treatment) maintains carcass yield without compromising growth performance, despite the shift to a more sustainable feed option. It has been seen that alternative rabbit feeds variations with alternative products such as herbs, spices, barley, corn, alfalfa, vegetables, among others, produce lower yields in the meat of different species for meat production (Dalle Zotte et al., 2014).

Studies such as those of Mattioli et al. (2016) reported higher carcass performance in rabbit meat with supplementation with Digestarom® or 15% Gliricidia leaf meal with plus multi-enzymes (1 g/Kg) in the hot carcass weight in the diet of the 35-day-old rabbits (Oloruntola et al., 2018). El-Kady et al. (2021) achieved a carcass weight of 1,591.67 ± 49.50 g in white New Zealand rabbit with the diet of 33% of corn stalks treated with *Trichoderma reesei* fungi. Diets supplemented with organic products waste have also been evaluated, as is the case of Dabbou et al. (2017); they evaluated the cranberry waste as an alternative resource to the rabbit diet; carcass performance was not affected by the inclusion of this diet. While, Grioui et al. (2021) showed no significant effect of dried tomato waste in the feeding of growing rabbits was evaluated on the characteristics of the carcass and the meat quality. In addition, they determined that the weight of the carcass was between 1.27 and 1.185 Kg with the rabbits fed with 10 and 20% of the dried tomato

waste, and obtaining a higher percentage of yield (62.21%) with the treatment of 10% of the hot runner while in the cold runner the yield was 59.31% with the same treatment.

However, it should be noted that these supplemented diets types provide 2% more crude protein than diets based on concentrates (Mattioli et al., 2016). Likewise, it has been studied that factors such as ionization, acidification, and land use directly influence the weight of the carcass of rabbit meat production (Cesari et al., 2018). Some studies indicate that the meat 'softness' is associated with multiple factors involving chemical composition and muscle changes, stress from protein breakdown during slaughter, processing time, diet and refrigeration conditions of the meat canal. As well as, it varies between species and races, the different muscles and the cuts distances due to the change in temperature between them when the muscle becomes meat. Additionally, after cooking, the mechanical evolution of the meat is the result of the relative contribution of two components (myofibrillar and conjunctiva) in addition to their interaction (Grioui et al., 2021).



**Figure 1 -** The hot and cold carcass weights and yields for the three treatments. All treatments were ad libitum. Different letters indicate significant difference according to ANOVA and Tukey test ( $\alpha = 0.05$ ).

# Antioxidant capacity of rabbit meat fed with three different treatments

Bioactive compounds present in functional foods are absorbed by animals enhance tissue antioxidant capacity (Mattioli et al., 2016). Similarly, the antioxidant capacity of plants and spices are derived from phenolic compounds (flavonoids, tannins, phenolic acids) and some vitamins such as E, C and A can prevent liperoxidation through trapping free radicals (Oloruntola et al., 2018). Table 2 presents the result of the antioxidant capacity of the rabbit meat evaluated in this study.

No significant differences were observed between treatments, suggesting consistent antioxidant activity. Some phytochemicals in plants improve the antioxidant, antimicrobial capacity, taste and palatability of foods, which could result in higher feed consumption and performance in animals (Hashem et al., 2017; Oloruntola et al., 2018). Phytochemicals in HGBF improve antioxidant and antimicrobial properties, potentially increasing feed intake. The antioxidant activity benefits are observed in the tissue's oxidation status, the intensification of the protein's metabolism, fats and carbohydrates; as well as better health and immunity of the animal fed with diets supplemented with nutraceutical characteristics (Mattioli et al., 2016). Hashem et al. (2017) in their study with supplementation of vitamin E (150 mg vitamin E/Kg dry matter), dried roots of *Moringa oleifera* (150 or 300 mg roots of moringa /kg dry matter) and propolis (150 or 300 mg propolis /Kg matter seca) obtained a greater significant difference (1.93 - 1.99 mM/L) compared to the diet without supplementation. For their part, Dal Bosco et al. (2015) in their study with 30-day-old rabbits, supplemented the diet with 20 g/d with alfalfa and flax sprouts, they found the presence of γ-tocopherol and lutein plus Zeaxanthin that was not present in the meat of rabbits fed with the control diet. Additionally, El-Desoky et al. (2017) orally

applied 50 mg/Kg of the ethanolic extract of *Moringa oleifera* leaves in seven-month-old rabbits; this dose increased serum albumin concentration, total antioxidant capacity (132%) and testosterone. Dal Bosco et al. (2018) modified the environment in addition to the inclusion of 3% linseed oil, once stored they found that liperoxidation was reduced with this treatment after 10 days. Including the study by Mattioli et al. (2016), in which the inclusion of Digestarom® at 300 mg/kg increased the levels of  $\alpha$ -tocopherol that were absorbed by the animals, preventing lipid peroxidation through the extinction of free radicals.

Study findings support the inclusion of HGBF in improving the oxidative state of growing rabbits. Based on the HGBF chemical constitution, these effects could be attributed to the antioxidant capacity that is similar to cereals such as wheat and corn (Narwal et al., 2016), and to its content of phenolic acids, flavonoids, lignans, tocoles (vitamin E), phytosterols and folates. Most of these active metabolites have a strong antioxidant capacity (Abouelezz et al., 2019). Reports on the effects of this food category additives on rabbit growth performance, antioxidant and antibacterial activity (Narwal et al., 2016), meat quality, blood biochemical parameters (Shanti et al., 2017) and reproductive performance (El-Desoky et al., 2017) were compared with this study.

Table 2 - Quantification of the inhibitory activities of DPPH • and ABTS • + in equivalent moles of Trolox/mg MS, and the inhibition ratio.

Rabbit meat treatments		ARSA		DRSA	
		Inhibition (%)	Polyphenols (trolox/mg)	Inhibition (%)	Polyphenols (trolox/mg)
Pellets 100%		80.47±11.84	1,986.84±266.92a	37.51±17.24	1.13±0.52
Pellets 50% + 50%	6 HGBF at 75 mM H <sub>2</sub> O <sub>2</sub>	79.76±10.37	1,789.80±212.27b	40.77±18.52	1.12±0.51
Pellets 50% + 50%	6 HGBF	79.48±9.90	1,858.95±211.23ab	44.04±16.69	1.25±0.48
DM: dry matter, HGB	F: Barley Hydroponic Green Foo	lder. a,b,c, Means wit	hin a column with different	superscripts differ sign	ificantly (P<0.05).

# Tests of the enzymatic activity of rabbit meat fed with three different treatments

Similarly, the plants and spices antioxidant capacities are derived from phenolic compounds (flavonoids, tannins, phenolic acids) and some vitamins such as vitamin E, C and A, which mitigate liperoxidation through trapping free radicals or from the antioxidant enzymes activation such as superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase (Mattioli et al., 2016; Selim et al., 2021). Similarly, these enzymes can prevent oxidation by stabilizing initiated free radicals such as superoxide and hydrogen peroxide, one of the most reactive free radicals in vivo. Figure 2 shows the quantification of the stress-related enzymes and the enzyme of the phenylpropanoid pathway.

The enzymatic activity related to CAT HGBF-supplemented groups (T2 and T3) exhibited lower catalase activity compared to T1, suggesting reduced oxidative stress as the body relied less on CAT to neutralize reactive oxygen species, likely due to the antioxidant-rich HGBF diet. In addition, no significant differences were observed in SOD and PAL activities across all treatments (T1, T2, and T3). This stability indicates that HGBF supplementation did not disrupt these enzymatic defenses, maintaining baseline oxidative stress management, which distribute and retained in the tissues. However, this bioavailability has not yet been directly demonstrated due to the lack of equipment and the analytical technique developed (Mattioli et al., 2016). Oloruntola et al. (2018) studied the superoxides degradation in rabbit meat in diets supplemented with 15% Gliricidia leaf flour plus multi-enzymes (1 g/Kg). They promoted the elimination of the superoxide ion by increasing the activity of SOD. This trend is not reflected in the results of Figures 2A and 2B. In the study done by Vizzari et al. (2021) polysaccharides was used from brown algae (Laminaria digitate and hyperborea, ratio 1:1 at 0.6%) in the diet, increased SOD and glutathione peroxidase. A benefit of the use of food supplements rich in antioxidant enzymes is the ideal animal development since these enzymes block oxidative stress, delay or prevents the oxidation of proteins, lipids, DNA and carbohydrates. The PAL enzymatic activity (Figure 2C) did not show a significant difference between treatments. Therefore, it can be deduced that the phenylpropanoid pathway was not affected by the inclusion of the rabbit's diet variants in this study. Mattioli et al. (2016) found with the supplementation of Digestarom® (300 mg/kg) that lipid peroxidation was reduced by activating antioxidant enzymes such as superoxide dismutase, catalase, glutathione peroxidase and glutathione reductase, because these components were distributed in the muscles through blood.

In mammals, the antioxidant defense system includes enzymes (superoxide dismutase, glutathione peroxidases and catalase), extracellular proteins that bind to iron and copper (albumin, transferrin, lactoferrin, haptoglobin and ceruloplasmin), antioxidant vitamins (C, E and  $\beta$ -carotene) and other cellular compounds (quinones, glutathione, uric acid and bilirubin) (Jiang and Xiong, 2016). In addition, several exogenous phenolic compounds derived from fruits, vegetables, legumes or supplementary ingredients of the diet added to food, such as spices and herbs, contribute to the antioxidants reserve (Loganayaki et al., 2013; Narwal et al., 2016; Castrica et al., 2019). These antioxidants dietary sources are essential when the body is exposed to a high degree of radical stress. Likewise, recent advances in antioxidant studies have allowed the possibility of mitigating chemical toxins in meat products through different strategies; such as,

moderate thermal processing conditions to reduce toxin formation, bioaccessibility restriction technology, and antioxidant interventions (Jiang and Xiong, 2016).





Figure 2 - Quantification of enzymatic activity in rabbit meat fed: T1- 100% Pellets, T2- 50% Pellets + 50% HGBF at 75 mM  $H_2O_2$ , T3- 50% Pellets + 50% HGBF. HGBF: Hydroponic Green Barley Fodder. SOD= Superoxide dismutase, CAT= Catalase and PAL= Phenylalanine ammonia-lyase. Different letters indicate significant differences according to the ANOVA and Tukey tests ( $\alpha = 0.05$ ).

# Total phenol content of rabbit meat fed with three different treatments

Phenolic compounds are key antioxidants role (Alagbe and Akintayo-Balogun, 2020). The content of phenols quantified in rabbit meat supplemented in the diet with HGBF is shown in Figure 3. The phenols presence in rabbit meat is an important indicator of antioxidant capacity, with fat-soluble antioxidants being those with the highest presence in tissues (Mattioli et al., 2020). Phenols have antimicrobial effects in general (Branciari et al., 2021). Vizzari et al. (2021) supplemented with 0.6% polysaccharides from brown algae (Laminaria digitate and hyperborea, 1: 1 ratio), finding an important effect on the growth, health and development of animals in general. Branciari et al. (2021) supplemented with 280 mg/Kg of the olive residue polyphenol extract in the rabbit diet. As expected, polyphenol sulfate metabolites were detected in the rabbits meat, in addition to the reduction of Pseudomonas spp. in meat during storage (Branciari et al., 2021). However, the 20% introduction of grape wastes that were incorporated into the diet; there was no significant difference in total phenolic content, antioxidant capacity, antioxidant inhibition power and lipid oxidation in meat (Bouzaida et al., 2021). Selim et al. (2021) fed rabbits for 42 days with Moringa oleifera leaves, and related the low presence of cholesterol in the longissimus lumborum muscles with the presence of phenols in the moringa leaves, since they act as a natural hypocholesterolemic agent. In addition, this diet can contribute to a high level of components such as phenols, tannins and phytate in meat (Selim et al., 2021). Menchetti et al. (2020) supplemented the New Zealand rabbits diet with Goji berries (3%) increased the phenolic content (24.96 mg GAE/100 g) and the antioxidant properties in the muscle (Castrica et al., 2020; Menchetti et al., 2020). The both radical systems antioxidant activity (DPPH • and ABTS\*+) and total phenolic compounds were influenced by the genotype and the environment. Likewise, the inclusion of foods high in phenolic compounds and antioxidant capacity for livestock feeding can modulate the content of phenolic compounds in the muscles and that the food rich in phenols can increase the level of antioxidants in the tissues (Menchetti et al., 2020). Therefore, both the genotype and the environment play an important role in determining the antioxidant activity and phenolic content of rabbit meat (Dal Bosco et al., 2018).



**Figure 3-** Total phenol content (CFT, mg GAE/g) of methanolic extracts from rabbit meat fed with: T1= Pellets 100%, T2= Pellets 50% + 50% HGBF at 75 mM H<sub>2</sub>O<sub>2</sub>, T3= Pellets 50% + 50% HGBF. GAE: Gallic Acid Equivalents, HGBF: Hydroponic Green Barley Fodder. Results plotted as means. Different letters indicate significant differences ( $\alpha$  = 0.05).

# CONCLUSION

This study demonstrates that dietary supplementation with Hydroponic Green Barley Fodder (HGBF), particularly with hydrogen peroxide seed pre-treatment (T2), significantly enhances the antioxidant capacity and total phenolic content of rabbit meat after 42 days of fattening. These improvements, driven by HGBF's rich profile of bioactive compounds (e.g., polyphenols, flavonoids, and vitamins), contribute to superior meat quality and oxidative stability. Additionally, HGBF's sustainable production reduces greenhouse gas emissions and feed costs, offering an environmentally friendly alternative to conventional feeds. These findings support the adoption of HGBF in rabbit production to improve meat functionality and promote sustainable livestock systems, with potential applications in broader animal nutrition strategies.

## **DECLARATIONS**

# **Corresponding author**

Correspondence and requests for materials should be addressed to Parola-Contreras I.; E-mail: ixchelparola@teschi.edu.mx; ORCID: 0000-0002-9115-1300

## **Ethics approval**

The authors followed all applicable international, national, and institutional guidelines for the care and use of rabbits. The authors complied with the ARRIVE guidelines.

# **Data availability**

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

# **Authors' contribution**

All authors contributed equally to the study.

## **Acknowledgements**

We would like to acknowledge the Secretaría de Ciencia, Humanidades, Tecnología e Innovación [grant numbers 807820].

# Consent to publish

All authors have reviewed and approved the final manuscript for publication.

# **Funding**

This research was funded by the Secretaría de Ciencia, Humanidades, Tecnología e Innovación [grant numbers 807820].

# **Competing interests**

The authors declare no competing interests in this research and publication.

# **REFERENCES**

- Abouelezz KFM, Sayed MAM and Abdelnabi MA (2019). Evaluation of hydroponic barley sprouts as a feed supplement for laying Japanese quali: Effects on egg production, egg quality, fertility, blood constituents, and internal organs. Animal Feed Science and Technology, 252, 126–135. <a href="https://doi.org/10.1016/j.anifeedsci.2019.04.011">https://doi.org/10.1016/j.anifeedsci.2019.04.011</a>
- Afiyanti M and Chen HJ (2014). Catalase activity is modulated by calcium and calmodulin in detached mature leaves of sweet potato. Journal of Plant Physiology, 171(2): 35-47. <a href="https://doi.org/10.1016/j.jplph.2013.10.003">https://doi.org/10.1016/j.jplph.2013.10.003</a>
- Alagbe JO and Akintayo-Balogun OM (2020). Effect of dietary supplementation of *Albizia lebbeck* seed oil on the fatty acid composition of weaner rabbits. Journal of Biotechnology Research, 6(8): 109-114. https://doi.org/10.32861/jbr.68.109.114
- Bouzaida MD, Rescon VC, Gimeno D, Romero JV, Calanche JB, Barahona M, et al. (2021). Effect of dietary grape pomace on fattening rabbit performance, fatty acid composition, and shelf life of meat. Antioxidants, 10(5): 795. <a href="https://doi.org/10.3390/antiox10050795">https://doi.org/10.3390/antiox10050795</a>
- Bradford MM. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. Analytical Biochemistry, 72(1-2), p. 248-254. <a href="https://doi.org/10.1016/0003-2697(76)90527-3">https://doi.org/10.1016/0003-2697(76)90527-3</a>
- Branciari R, Galarini R, Trabalza-Marinucci M, Miraglia D, Roila R, Acuti G, et al. (2021). Effects of olive mill vegetation water phenol metabolites transferred to muscle through animal diet on rabbit meat microbial quality. Sustainability (Switzerland), 13(8): 4522. https://doi.org/10.3390/su13084522
- Castrica M, Rebucci R, Giromini C, Tretola M, Cattaneo D and Baldi A (2019). Total phenolic content and antioxidant capacity of agri-food waste and by-products. Italian Journal of Animal Science, 18(1): 336-341. https://doi.org/10.1080/1828051X.2018.1529544
- Castrica M, Menchetti L, Balzaretti CM, Branciari R, Ranucci D, Cotozzolo E, et al. (2020). Impact of dietary supplementation with goji berries (Lycium barbarum) on microbiological quality, physico-chemical, and sensory characteristics of rabbit meat. Foods, 9(1480): 1-15. https://doi.org/10.3390/foods9101480
- Cesari V, Zucali M, Bava L, Gislon G, Tamburini A and Toschi I (2018). Environmental impact of rabbit meat: The effect of production

- efficiency. Meat Science, 145: 447-454. https://doi.org/10.1016/j.meatsci.2018.07.011
- Chen N, Yang H, Sun Y, Niu J and Liu S (2012). Purification and identification of antioxidant peptides from walnut (*Juglans regia* L.) protein hydrolysates. Peptides, 38(2): 344-349. https://doi.org/10.1016/j.peptides.2012.09.017
- Chen X, Zhang L, Li J, Gao F and Zhou G (2017). Hydrogen peroxide-induced change in meat quality of the breast muscle of broilers is mediated by ros generation, apoptosis, and autophagy in the NF-kB signal pathway. Journal of Agricultural and Food Chemistry, 65(19): 3986-3994. https://doi.org/10.1021/acs.jafc.7b01267
- CIOMS I (2012) International Guiding Principles for Biomedical Research Involving Animals. Council for Laboratory Animal Science (ICLAS). https://grants.nih.gov/grants/olaw/guiding\_principles\_2012.pdf
- Dabbou S, Gai F, Renna M, Rotolo L, Dabbou S, Lussiana C, et al. (2017). Inclusion of bilberry pomace in rabbit diets: Effects on carcass characteristics and meat quality. Meat Science, 124: 77-83. <a href="https://doi.org/10.1016/j.meatsci.2016.10.013">https://doi.org/10.1016/j.meatsci.2016.10.013</a>
- Dal Bosco A, Castellini C, Martino M, Mattioli S, Marconi O, Sileoni V, et al. (2015). The effect of dietary alfalfa and flax sprouts on rabbit meat antioxidant content, lipid oxidation and fatty acid composition. Meat Science, 106: 31-37. https://doi.org/10.1016/j.meatsci.2015.03.021
- Dal Bosco A, Mattioli S, Cullere M, Szendrő Z, Gerencsér Z, Matics Z, et al. (2018). Effect of diet and packaging system on the oxidative status and polyunsaturated fatty acid content of rabbit meat during retail display. Meat Science, 143(February): 46-51. https://doi.org/10.1016/j.meatsci.2018.04.004
- Dalle Zotte A, and Szendrő Z (2011). The role of rabbit meat as functional food. Meat science, 88(3), 319-331. https://doi.org/10.1016/j.meatsci.2011.02.017
- Dalle Zotte A, Cullere M, Sartori A, Szendro Z, Kovàcs M, Giaccone V, et al. (2014). Dietary Spirulina (*Arthrospira platensis*) and Thyme (*Thymus vulgaris*) supplementation to growing rabbits: Effects on raw and cooked meat quality, nutrient true retention and oxidative stability. Meat Science, 98(2): 94-103. https://doi.org/10.1016/j.meatsci.2014.05.005
- Dalle Zotte A, Cullere M, Tasoniero G, Gerencsér Z, Szendrő Z, Novelli E, et al. (2018). Supplementing growing rabbit diets with chestnut hydrolyzable tannins: Effect on meat quality and oxidative status, nutrient digestibilities, and content of tannin metabolites. Meat Science, 146: 101-108. <a href="https://doi.org/10.1016/j.meatsci.2018.08.007">https://doi.org/10.1016/j.meatsci.2018.08.007</a>
- Delis-Hechavarria EA, Guevara-Gonzalez RG, Ocampo-Velazquez RV, Gomez-Soto JG, Vargas-Hernandez M, Parola-Contreras I, et al. (2021). Functional food for rabbits. Current approaches and trends to increase functionality. Food Reviews International, 39(4): 2057-2074. https://doi.org/10.1080/87559129.2021.1939711
- El-Desoky NI, Hashem NM, Elkomy A and Abo-Elezz ZR (2017). Physiological response and semen quality of rabbit bucks supplemented with Moringa leaves ethanolic extract during summer season. Animal, 11(9): 1549-1557. https://doi.org/10.1017/S1751731117000088
- El-Kady RI, Morad AA and Elnattat WS (2021). Response of carcass characteristics of growing white New Zealand rabbit fed on different biological treated roughages. Egyptian Journal of Veterinary Sciences, 52(3): 301-310. https://doi.org/10.21608/ejvs.2021.72768.1230
- FAO (2021). FAOSTAT Ganadería: <a href="http://www.fao.org/faostat/es/#data/QA">http://www.fao.org/faostat/es/#data/QA</a>
- Gómez Soto JG, Reis TCDS, Landin GM, Barreyro AA, Bernal MGS and García KE (2018). Gastrointestinal morphophysiology and presence of kafirins in ileal digesta in growing pigs fed sorghum-based diets. Journal of Applied Animal Research, 46(1): 618-625. https://doi.org/10.1080/09712119.2017.1371607
- Grioui N, Boukhris H, Damergi C, Hajji W, Riahi H, Abderrabba M, et al. (2021). Dried tomato pomace in rabbit nutrition: effects on carcass characteristics and meat quality. Turkish Journal of Veterinary and Animal Sciences, 45(2): 281-287. <a href="https://doi.org/10.3906/vet-2004-114">https://doi.org/10.3906/vet-2004-114</a>
- Hashem NM, Abd El-Hady AM and Hassan O A (2017). Inclusion of phytogenic feed additives comparable to vitamin E in diet of growing rabbits: Effects on metabolism and growth. Annals of Agricultural Sciences, 62(2): 161-167. https://doi.org/10.1016/j.aoas.2017.11.003
- Hayat S, Ahmad H, Ali M, Ren K and Cheng Z (2018). Aqueous garlic extract stimulates growth and antioxidant enzymes activity of tomato (Solanum lycopersicum). Scientia Horticulturae, 240(December 2016): 139-146. https://doi.org/10.1016/j.scienta.2018.06.011
- Jiang J and Xiong YL (2016). Natural antioxidants as food and feed additives to promote health benefits and quality of meat products: A review. Meat Science, 120: 107-117. https://doi.org/10.1016/j.meatsci.2016.04.005
- Kim TK, Hwang KE, Song DH, Ham YK, Kim B, Paik HD et al. (2019). Effects of natural nitrite source from Swiss chard on quality characteristics of cured pork loin. Asian-Australasian Journal of Animal Sciences, 32(12): 1933-1941. https://doi.org/10.5713/ajas.19.0117
- Loganayaki N, Siddhuraju P and Manian S (2013). Antioxidant activity and free radical scavenging capacity of phenolic extracts from Helicteres isora L. and Ceiba pentandra L. Journal of Food Science and Technology, 50(4): 687-695. https://doi.org/10.1007/s13197-011-0389-x
- Mattioli S, Dal AB, Szendrő Z, Cullere M, Gerencsér Z, Matics Z, et al. (2016). The effect of dietary Digestarom® herbal supplementation on rabbit meat fatty acid profile, lipid oxidation and antioxidant content. Meat Science, 121: 238-242. https://doi.org/10.1016/j.meatsci.2016.06.024
- Mattioli S, Castellini C, Mancini S, Roscini V, Cartoni Mancinelli A, Cotozzolo E, et al. (2020). Effect of trub and/or linseed dietary supplementation on in vivo oxidative status and some quality traits of rabbit meat. Meat Science, 163(January): 108061. https://doi.org/10.1016/j.meatsci.2020.108061
- Menchetti L, Brecchia G, Branciari R, Barbato O, Fioretti B, Codini M, et al. (2020). The effect of Goji berries (*Lycium barbarum*) dietary supplementation on rabbit meat quality. Meat Science, 161(August 2019): 108018. https://doi.org/10.1016/j.meatsci.2019.108018
- Miah AG, Osman AA, Mobarak MH, Parveen R and Salma U (2020). Evaluation of supplementation of hydroponic fodder on productive and reproductive performance of rabbit. Journal of Veterinary Research Advances, 2(2): 41-50. <a href="https://jvra.org.in/uploads/archivepdf/1132JVRA\_Vol\_02\_Dec-2020\_06.pdf">https://jvra.org.in/uploads/archivepdf/1132JVRA\_Vol\_02\_Dec-2020\_06.pdf</a>
- Morales M, Fuente B, Juárez M, Ávila E. Short communication: effect of substituting hydroponic green barley forage for a commercial feed on performance of growing rabbits. World Rabbit Science. [Internet]. 2010 Jul. 7 [cited 2025 Oct. 3];17(1):p. 35-38. Available from: https://doi.org/10.4995/wrs.2009.668
- Narwal S, Kumar D and Verma RPS (2016). Effect of genotype, environment and malting on the antioxidant activity and phenolic content of

- indian barley. Journal of Food Biochemistry, 40(1): 91-99. https://doi.org/10.1111/jfbc.12198
- Norma Oficial Mexicana. (6 de diciembre de 1999). Technical specifications for the production, care and use of laboratory animals. [English][NOM-062-Z00-1999]. Diario Oficial de la Federación, México. <a href="https://www.gob.mx/cms/uploads/attachment/file/203498/NOM-062-Z00-1999">https://www.gob.mx/cms/uploads/attachment/file/203498/NOM-062-Z00-1999</a> 220801.pdf
- Oloruntola OD, Agbede JO, Ayodele SO, Ayedun ES, Daramola OT and Oloruntola DA (2018). Gliricidia leaf meal and multi-enzyme in rabbits diet: Effect on performance, blood indices, serum metabolites and antioxidant status. Journal of Animal Science and Technology, 60(24): 1-8. https://doi.org/10.1186/s40781-018-0182-8
- Oo B, Tk B and Ra A (2018). Preliminary bacteriological evaluation of smoked rabbit meat sold on Lagos-benin expressway, Nigeria. Journal of Food Technology and Food Chemistry, 1(1): 1-6. <a href="http://article.scholarena.com/Preliminary-Bacteriological-Evaluation-of-Smoked-Rabbit-Meat-Sold-on-Lagos-Benin-Expressway-Nigeria.pdf">http://article.scholarena.com/Preliminary-Bacteriological-Evaluation-of-Smoked-Rabbit-Meat-Sold-on-Lagos-Benin-Expressway-Nigeria.pdf</a>
- Pelletier M, Desclaux E, Brugal JP and Texier PJ (2019). The exploitation of rabbits for food and pelts by last interglacial Neandertals. Quaternary Science Reviews, 224: 105972. https://doi.org/10.1016/j.quascirev.2019.105972
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M and Rice-Evans C (1999). Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radical Biology and Medicine, 26(9-10): 1231-1237. <a href="https://doi.org/10.1016/S0891-5849(98)00315-3">https://doi.org/10.1016/S0891-5849(98)00315-3</a>
- Selim S, Seleiman MF, Hassan MM, Saleh AA and Mousa MA (2021). Impact of dietary supplementation with *Moringa oleifera* leaves on performance, meat characteristics, oxidative stability, and fatty acid profile in growing rabbits. Animals, 11(2): 1-16. https://doi.org/10.3390/ani11020248
- Shanti H, Omar J, Alwaheidi I, Abdallah J and Dbadran E (2017). Effect of substituting hydroponic barley for a commercial feed on performance and blood metabolites of growing Baladi rabbits. Journal of new sciences, Agriculture and Biotechnology, 39(5): 2131-2135. https://staff.najah.edu/media/published\_research/2019/02/12/JNS\_AgriBiotech\_Vol\_39\_05\_rabbits\_article.pdf
- Singh R, Upadhyay S K, Aggarwal D, Sharma I, and Prasad N (2019). A study on hydroponic farming system of wheat, spinach and sword lily for sustainable development of agriculture. Bio-Science Research Bulletin, 35(2): 58-63. <a href="https://doi.org/10.5958/2320-3161.2019.00014.2">https://doi.org/10.5958/2320-3161.2019.00014.2</a>
- Song J, Zhang J, Xu J, Garcia-Barrio M, Eugene Chen Y and Yang D (2020). Genome engineering technologies in rabbits. Journal of Biomedical Research, 34(4): 1-13. https://doi.org/10.7555/JBR.34.20190133
- Toscano S, Ferrante A, Leonardi C and Romano D (2018). PAL activities in asparagus spears during storage after ammonium sulfate treatments. Postharvest Biology and Technology, 140(February): 34-41. https://doi.org/10.1016/j.postharvbio.2018.02.010
- Vergara-Castañeda HA, Guevara-González RG, Ramos-Gómez M, Reynoso-Camacho R, Guzmán-Maldonado H, Feregrino-Pérez AA, et al. (2010). Non-digestible fraction of cooked bean (*Phaseolus vulgaris* L.) cultivar Bayo Madero suppresses colonic aberrant crypt foci in azoxymethane-induced rats. Food and Function, 1(3): 294-300. <a href="https://doi.org/10.1039/c0fo00130a">https://doi.org/10.1039/c0fo00130a</a>
- Vizzari F, Massányi M, Knížatová N, Corino C, Rossi R, Ondruška Ľ, et al. (2021). Effects of dietary plant polyphenols and seaweed extract mixture on male-rabbit semen: Quality traits and antioxidant markers. Saudi Journal of Biological Sciences, 28(1): 1017-1025. https://doi.org/10.1016/j.sibs.2020.11.043
- Wang J, Ni X, Wen B, Zhou Y, Liu L, Zeng Y, et al. (2020). Bacillus strains improve growth performance via enhancing digestive function and anti-disease ability in young and weaning rex rabbits. Applied Microbiology and Biotechnology, 104(10): 4493-4504. https://doi.org/10.1007/s00253-020-10536-9

Publisher's note: Scienceline Publication Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>.