








EFFECT OF EMULSIFIED HIGH-PUFA OILS ENRICHED WITH NATURAL ANTIOXIDANTS ON IN VITRO GOAT RUMEN FERMENTATION PARAMETERS AND NUTRIENT DEGRADABILITY

Dilla Mareistia FASSAH¹✉ , Manik Nurul ALFIYYAH¹ , Safiena Maura AZKA¹ , Shafira Pramesti DAMAYANTI¹ , Yusti PUJIAWATI^{1,2} , Lilis KHOTIJAH¹ , and Novia Amalia SHOLEHA³ 

¹Department of Animal Nutrition and Feed Technology, Faculty of Animal Science, IPB University, Bogor, 16680, Indonesia

²Research Center for Animal Husbandry, Research Organization for Agriculture and Food, National Research and Innovation Agency Indonesia, Cibinong Science Center, Jalan Raya Jakarta-Bogor, Cibinong, Bogor, 16915, Indonesia

³College of Vocational Studies, IPB University, Jalan Kumbang No. 14, Bogor, 16151, Indonesia

✉ Email: dilla_mareistia@apps.ipb.ac.id

Supporting Information



ABSTRACT: This study aims to evaluate the effect of emulsified high-PUFA oils enriched with natural antioxidants derived from jackfruit leaf extract on in vitro goat rumen fermentability and degradability. The study used a completely randomized design with two treatments and 5 replications. The treatments were supplementation with high-PUFA oils + natural antioxidants (P1, as the control), and supplementation with emulsified high-PUFA oils + natural antioxidants (P2). The parameters observed were rumen pH, total volatile fatty acid (VFA) concentration, NH₃ concentration, bacterial population, protozoa population, dry matter digestibility (DMD), and organic matter digestibility (OMD). The results showed that supplementing goat feed ration with emulsified high-PUFA oils enriched with natural antioxidants did not alter ($P > 0.05$) rumen pH, total VFA, NH₃ concentration, bacterial and protozoal population, DMD, or OMD. Overall, emulsified high-PUFA oils with natural antioxidants did not alter the examined rumen fermentation parameters and degradability. Thus, emulsified high-PUFA oils with natural antioxidants can be potentially used as an effective feed supplement to support goats' performance.

Keywords: Emulsification, Feed supplement, Fermentation characteristics, Jackfruit leaves, Polyunsaturated fatty acid.

RESEARCH ARTICLE
 PII: S222877012600014-16
 Received: September 11, 2025
 Revised: March 28, 2026
 Accepted: March 29, 2026

INTRODUCTION

Previous studies have shown that polyunsaturated fatty acids (PUFA), particularly n-3 and n-6, have a positive effect on the reproductive function, health, and performance of goats (Almeida et al., 2019; Geng et al., 2020). However, ruminants cannot synthesize n-3 and n-6 PUFAs de novo, so the fatty acids must be supplied through their diet (Roque-Jiménez et al., 2021). Flaxseed and sunflower oils are notable sources of essential n-3 and n-6 PUFAs. Flaxseed oil is rich in n-3 fatty acids, contains about 50% α -linoleic acid (ALA), which can be metabolized into eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), and has anti-inflammatory and antiatherosclerosis effects (Liao et al., 2021; Romanić et al., 2021). Sunflower seed oil is predominantly composed of n-6 PUFAs, particularly linoleic acid (LA), which reaches 60%-75% (Akkaya, 2018). However, high-PUFA oil supplementation in ruminant diets may negatively affect rumen fermentation by inhibiting microbial activity, particularly fiber-degrading bacteria, thereby reducing fermentation efficiency when supplied at inappropriate levels or in unprotected forms (Mirzaei-Alamouti et al., 2021). A previous study has also reported that dietary lipid supplementation exceeding 4-6% of dry matter may negatively affect rumen fermentation (Ribeiro et al., 2025).

In addition, PUFA metabolism may induce oxidative stress in livestock, leading to increased reactive oxygen species (ROS) and altering PUFAs' biological functions. Exogenous antioxidants can be used to prevent the side effects of PUFA metabolism (Fassah and Khotijah, 2016). A natural antioxidant derived from Jackfruit leaf extract, containing 0.34% antioxidant compounds, has been proposed as a strategy to mitigate oxidative stress (Pujiawati et al., 2023). Ojwang et al. (2017) reported that jackfruit leaves contain various bioactive compounds, including flavonoids (5.02-6.70 mg/g), tannins (1.97-2.56 mg/g), and phenolic compounds (30.92 - 37.39 mg/g). Nevertheless, directly mixing PUFA oils with jackfruit leaf extract may pose problems, as differences in polarity between hydrophilic antioxidants and lipophilic PUFA oils result in poor mixing efficiency, ineffective stabilization, and limited protective effects (Roschel et al., 2019).

Emulsification has emerged as a promising approach to improve the physical stability and dispersion of PUFA oils and antioxidants. Emulsification allows for a more homogeneous mixture of PUFA oils and antioxidants by adding surfactants that reduce interfacial tension between two liquid phases (Saxena and Mandal, 2022). Tween 80 is a non-ionic surfactant commonly used in emulsions that generally do not alter the pH since it does not dissociate into ions that could affect pH levels (López-Martínez and Rocha-Urbe, 2018; Pernin et al., 2019). Therefore, emulsifying high-PUFA oils with natural antioxidants may improve ruminal fermentation efficiency by stabilizing high-PUFA oils and natural

antioxidants without impairing nutrient degradability. However, studies on the impact of emulsified high-PUFA oils-antioxidants combinations on rumen fermentation remain limited. Therefore, this study aims to evaluate the effect of emulsified high-PUFA oils enriched with natural antioxidants on in vitro rumen fermentation and degradability in goats.

MATERIALS AND METHODS

The research was conducted from October to December 2024 at the Meat and Draught Animal Nutrition Laboratory, IPB University, Indonesia. The in vitro analysis using goat rumen fluid was approved by the Animal Ethics Committee of the Directorate of Research and Innovation Licensing Governance and Scientific Authority (No. 123/KE.02/SK/06/2023).

Antioxidant extraction

The antioxidant compounds were extracted from jackfruit leaves using a maceration method in 70% ethanol (Najini and Wahdaningsih, 2024). The jackfruit leaves were briefly oven-dried at 60°C, then ground into a fine powder. A total of 1 kg of jackfruit leaves powder was macerated in 70% ethanol, then kept at room temperature for 24 hours. The mixture was then filtered using filter paper and concentrated using a rotary evaporator to form a viscous extract.

Feed supplement preparation

The feed supplement was prepared in oil-in-water (O/W) emulsion containing 15% high-PUFA oils (flaxseed oil and sunflower oil), 10% Tween 80, and 75% distilled water with 15 mg of antioxidant compounds derived from jackfruit leaf extracts. The oil phase consisted of 62% sunflower oil and 38% flaxseed oil (w/w), chosen to balance n-6/n-3 PUFA profile. Then, the mixture was homogenized using a sonicator (Lab Ultrasonic Homogenizer, Tefic Biotech Co., China) at 60 °C, power 100W, and 30 minutes of sonication to ensure a uniform distribution of the oil phase in the water phase (oil-in-water) (Chen et al., 2023). For both treatments, the total amount of high-PUFA oils and antioxidant compounds was standardized to deliver 150 mg of total oil and 15 mg of antioxidant per 100 mL emulsion.

In vitro ruminal fermentation characteristics

A Completely Randomized Design with 2 treatments and 5 replications per treatment was applied in this study. The treatments consisted of adding high-PUFA oil and natural antioxidant compounds separately to rumen fluid (P1; the control), or adding an emulsified high-PUFA oil enriched with natural antioxidant compounds to rumen fluid (P2). The treatments were evaluated using the in vitro two-stage method described by Tilley and Terry (1963). The formula and nutrient composition of the basal diet, as well as the composition of the supplement, are provided in Table 1.

Table 1 - Formula and nutrient compositions of the treatment feed rations

Feedstuff	Composition (%)	P1	P2
<i>Pennisetum purpureum</i>		30.00	30.00
Cassava waste		15.51	15.51
Copra meal		20.68	20.68
Soybean meal		10.34	10.34
Pollard		17.17	17.17
CaCO ₃		0.72	0.72
Premix		0.72	0.72
Salt		0.72	0.72
Molasses		4.14	4.14
Total		100	100
Supplement			
High-PUFA oils and natural antioxidants		2.31	-
Emulsified high-PUFA oils and natural antioxidants		-	2.31
Nutrient composition (%DM)			
Dry matter (DM) ¹		89.22	89.22
Ash ¹		7.48	7.48
Crude protein (CP) ¹		15.02	15.02
Ether extract (EE) ¹		2.01	2.01
Crude fiber (CF) ¹		15.13	15.13
Nitrogen-free extract (NFE) ¹		60.38	60.38
Total digestible nutrient (TDN) ²		69.14	69.14

P1: High-PUFA oils and natural antioxidants supplementation; P2: Emulsified High-PUFA oils and natural antioxidants supplementation; ¹Analyzed using Near infrared spectroscopy (NIRS); ²TDN = 40.265 - 1.379 (CF) + 1.1903 (EE) + 1.363 (NFE) + 0.1969 (CP) (Wardeh, 1981)

Rumen fluid was collected from three healthy donor goats using a stomach tube method before morning feeding (Shen et al., 2012). The animals were fed a basal diet consisting of 70% concentrate and 30% elephant grass (Crude protein = 9.52%, TDN = 66.45%). Rumen fluid was filtered through six-layer gauze, pooled into a single homogenized sample to reduce individual variation, and placed in an insulated bottle at 39 °C. A total of 0.5 g of the sample was incubated under anaerobic conditions at 39 °C with 10 mL of rumen fluid and 40 mL of McDougall buffer solution. Both supplements were added at a concentration of 23 mg/g sample. After 48 hours of incubation, 2-3 drops of HgCl₂ were added, followed by centrifugation at 3000 rpm for 15 minutes. The samples were filtered and separated into residue and liquid samples. The residue sample was collected, dried at 105 °C for 24 hours, and combusted in a muffle furnace for 4 hours (AOAC, 2019) to determine dry matter (DMD) and organic matter degradability (OMD). Meanwhile, the liquid sample was collected to analyze rumen pH, total volatile fatty acid (VFA), ammonia (NH₃), and microbial population. Although the method primarily assessed nutrient degradability, the rumen fermentation parameters were also determined at the end of incubation as endpoint indicators of rumen fermentation conditions.

The liquid samples were tested for rumen pH using a pH meter. The concentration of ammonia (NH₃) was calculated using the method by Chaney and Marbach (1962) using UV-Vis spectrophotometry at 630 nm. The total VFA was measured using a steam distillation method (Despal et al., 2022). The total culturable bacterial population in the rumen was determined using the spread plate method with Brain Heart Infusion (BHI) medium, as described by Oliveira et al. (2021). The rumen protozoa population was analyzed following Ogimoto and Imai (1981). Briefly, 2 drops of the rumen fluid sample were mixed with a trypan blue formalin saline solution in a 1:1 ratio, and the mixture was observed under a microscope.

Data analysis

The data obtained were analyzed using an independent sample t-test in the statistical software SPSS 25.0. Data was considered significantly different between treatments at P<0.05.

RESULTS AND DISCUSSION

Ruminal fermentation characteristics

The results showed that adding emulsified high-PUFA oils enriched with natural antioxidants to the ration did not significantly affect (P>0.05) pH value, total VFA, ammonia concentration, and bacterial and protozoal population (Table 2), indicating the emulsion's lack of adverse effect on rumen fermentation. In this study, the emulsion containing high-PUFA oils and natural antioxidants was stabilized with Tween 80, which has been shown to remain stable during digestion (Lamothe et al., 2019). Therefore, Tween 80 may have contributed to maintaining stable rumen pH. The ruminal pH values in this study were within the normal range of 6-7 (McDonald et al., 2022), which supports the growth and activity of rumen microbes required for optimal feed degradation. Therefore, the emulsion of high-PUFA oils-enriched with antioxidants did not interfere with the rumen's buffering capacity.

Table 2 – Rumen fermentation characteristics due to Emulsified high-PUFA oil with natural antioxidant supplementation

Parameters	Treatments	P1	P2	P-value
pH		7.10 ± 0.24	6.95 ± 0.30	0.523
Bacteria population (log cfu/mL)		6.25 ± 0.85	6.95 ± 0.20	0.446
Protozoa population (log cell/mL)		6.21 ± 0.12	6.19 ± 0.03	0.767
NH ₃ (mM)		8.07 ± 1.91	8.12 ± 0.69	0.971
Total VFA (mM)		100.04 ± 4.23	111.76 ± 6.41	0.057

P1 = high-PUFA oils and natural antioxidants supplementation, P2 = emulsified high-PUFA oils and natural antioxidants supplementation. VFA = volatile fatty acid, NH₃ = ammonia

Furthermore, the emulsified high-PUFA oils with natural antioxidants had no significant effect (P>0.05) on both the total ruminal bacteria and protozoa population. The lack of significant changes in the rumen microbial population may be due to the protective role of emulsifiers, which prevent oils and antioxidants from directly interacting with bacteria and protozoa. Previous study has reported that nano-emulsified oils modulate the proportions of volatile fatty acids in ruminal cultures, but do not affect total bacterial counts (El-Sherbiny et al., 2016). In addition, caffeic acid, a phenolic antioxidant, did not significantly affect bacterial counts when administered in emulsified form (Sørensen et al., 2008). In this study, the bacterial population findings align with those of Suharti et al. (2019), who reported that the total ruminal bacterial population with the addition of canola and flaxseed oil calcium soap ranged from 6.00 to 6.68 log cfu/mL. Meanwhile, the protozoa population in this study aligns with Zahera et al. (2024), who reported that the proportion of unprotected to protected fat supplementation in dairy rations ranged from 5.96 to 6.27 log cells/mL.

Next, the emulsified high-PUFA oils enriched with natural antioxidants did not change ($P>0.05$) NH_3 concentration. This finding indicates that protein degradation and microbial nitrogen assimilation remained unaffected by the supplementation form. In this study, the overall rumen microbial population remained unchanged with the addition of emulsified high-PUFA oils enriched with natural antioxidants, which may have maintained the ammonia concentration. Rumen-ammonia-producing bacteria, such as *Clostridium aminophilum*, are primarily responsible for producing ammonia from protein degradation (Adeniyi et al., 2023; Chanu et al., 2024). Additionally, protozoa such as *Entodinium caudatum* engulf bacteria and release oligopeptides and amino acids, which are fermented to ammonia by amino acid-fermenting bacteria (Park and Yu, 2023). The NH_3 concentration in this study aligns with that reported by Riestanti et al. (2024), who found that unsaturated fat supplementation in the form of calcium soap and prill fat resulted in NH_3 concentrations ranging from 7.73 to 9.73 mM. Therefore, our research indicates that supplementing goat feed rations with emulsified high-PUFA oils enriched with natural antioxidants can maintain normal NH_3 concentrations, supporting a well-functioning rumen.

Moreover, the total VFA was not altered by the supplementation of high-PUFA oils enriched with natural antioxidants (Table 2). Emulsification reduces the tension between oil and water, allowing better dispersion of oil droplets (Losada - Barreiro et al., 2020). In addition, the antioxidants at the oil-water interface effectively enhance the oxidative stability of emulsions (Li et al., 2022). The absence of significant changes in total VFA concentration suggests that the emulsified high-PUFA oils enriched with natural antioxidants neither stimulated nor suppressed overall fermentation activity. This result also indicates that emulsification may have played a protective role, maintaining PUFA and antioxidant function without affecting microbial fermentation. The total VFA levels in this study were comparable with those reported by Mukhtiani et al. (2024), ranging from 87.52 to 164.38 mM. Thus, emulsified high-PUFA oils with natural antioxidants may enhance supplementation efficiency in goats. However, further in vivo studies are required to confirm the protective effect of emulsification.

Ruminal in vitro degradability

Adding emulsified high-PUFA oils enriched with natural antioxidants did not influence ($P>0.05$) DMD and OMD (Table 3). The lack of a significant difference in DMD and OMD levels with supplementation in the emulsified form may be caused by the content of active compounds within the emulsion that did not interfere with the rumen microbial population and rumen pH value. The microbial population and pH conditions in the rumen may facilitate the fermentation and feed degradation process (Mickdam et al., 2016). In our study, the DMD and OMD remain unchanged as the emulsified substrates may not be fully utilized by rumen microbes (Eftaxias et al., 2021). Additionally, the antioxidant components in the emulsion may have created a barrier that limits the nutrient degradability (Obando et al., 2015). The digestibility of DM and OM in this study aligns with those reported by Reproto (2025), which ranged from 42.01% to 76.90% and from 65.1% to 88.36%, respectively. Thus, adding emulsified high-PUFA oils enriched with natural antioxidants to the goat ration results in consistent DMD and OMD levels, reflecting its potency as a feed supplement without altering rumen fermentation or degradability.

Table 3 - Ruminal dry matter and organic matter degradability

Parameters	Treatments		P-value
	P1	P2	
Dry matter degradability (DMD)	76.58 ± 4.73	74.97 ± 2.98	0.643
Organic matter degradability (OMD)	77.28 ± 6.26	74.80 ± 2.65	0.562

P1 = high-PUFA oils and natural antioxidants supplementation, P2 = emulsified high-PUFA oils and natural antioxidants supplementation

CONCLUSION

This study found that emulsified high-PUFA oils enriched with natural antioxidants does not alter rumen pH, microbial population, total VFA, ammonia concentration, and the degradability of dry matter and organic matter. Thus, emulsified high-PUFA oils enriched with natural antioxidants may be used as an efficient and safe feed supplement for goats. Nevertheless, further in vivo studies are strongly required to assess the long-term effects of supplementation on rumen fermentation and animal performance.

DECLARATIONS

Corresponding author

Correspondence and requests for materials should be addressed to Dilla Mareistia Fassah. E-mail: dilla_mareistia@apps.ipb.ac.id; ORCID: 0000-0003-4309-9218.

Data availability

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Authors' contribution

Dilla Mareistia Fassah: Conceptualization, methodology, data curation, data analysis, writing – original draft; Manik Nurul Alfiyyah; Methodology, data curation, data analysis; Safiena Maura Azka: Methodology, data curation, data analysis; S P Damayanti: Methodology, data curation, data analysis; Y Pujiawati: Conceptualization, methodology, writing – review and editing; Lilis Khotijah: Conceptualization, methodology, validation, writing – review and editing; N A Sholeha: Conceptualization, methodology.

Acknowledgements

The authors gratefully acknowledge the Directorate General of Higher Education, Science, and Technology, Ministry of Higher Education, Science, and Technology of the Republic of Indonesia, which supports this study through the BIMA Program 2024 grant (Contract No. 027/E5/PG.02.00.PL/2024).

Ethical approval

All the experimental procedures were conducted by the Animal Ethics Committee of the Directorate of Research and Innovation Licensing Governance and Scientific Authority (No 123/KE.02/SK/06/2023).

Funding

Directorate General of Higher Education, Science, and Technology, Ministry of Higher Education, Science, and Technology of the Republic of Indonesia provided financial support for this study through the BIMA Program 2024 grant (Contract No. 027/E5/PG.02.00.PL/2024).

Consent to publish

All authors agree to the publication of this manuscript.

Competing interests

The authors have not declared any competing interests.

REFERENCES

- Adeniyi A, Bello I, Mukaila T, Monono E and Hammed A (2023). Developing rumen mimicry process for biological ammonia synthesis. *Bioprocess and Biosystems Engineering*, 46(7): 1011–1020. <https://doi.org/10.1007/s00449-023-02880-7>
- Akkaya MR (2018). Fatty acid compositions of sunflowers (*Helianthus annuus* L.) grown in east mediterranean region. *Rivista Italiana Delle Sostanze Grasse*, 95: 239–247. <https://openaccess.atu.edu.tr/items/f679c88b-2d97-40ee-951b-4e648b05af56/full>
- Almeida OC, Ferraz MVC, Susin I, Gentil RS, Polizel DM, Ferreira EM, et al. (2019). Plasma and milk fatty acid profiles in goats fed diets supplemented with oils from soybean, linseed or fish. *Small Ruminant Research*, 170: 125–130. <https://doi.org/10.1016/j.smallrumres.2018.11.002>
- AOAC (2019). Association of Official Analytical Chemists. Official Methods of Analysis of AOAC International, 21st Edition. Association of Official Analytical Chemists International, Washington DC.
- Chaney AL and Marbach EP (1962). Modified reagents for determination of urea and ammonia. *Clinical Chemistry*, 8(2): 130–132. <https://doi.org/10.1093/clinchem/8.2.130>
- Chanu YM, Paul SS, Dey A and Andonissamy J (2024). Deciphering hyperammonia-producing bacteria (HAB) in the rumen of water buffaloes (*Bubalus bubalis*) and their inhibition through plant extracts and essential oils. *Microorganisms*, 12(10): 2040. <https://doi.org/10.3390/microorganisms12102040>
- Chen Y, Cheng H and Liang L (2023). Effect of oil type on spatial partition of resveratrol in the aqueous phase, the protein interface and the oil phase of O/W emulsions stabilized by whey protein and caseinate. *Antioxidants*, 12(3): 589. <https://doi.org/10.3390/antiox12030589>
- Despal D, Irmadani D, Permana IG, Zahera R and Nuraina N (2022). Effect of different unsaturated fatty acids sources on in-vitro fermentability and digestibility of ration in dairy cattle. *Online Journal of Animal and Feed Research*, 12(3): 154-159. <https://doi.org/10.51227/ojafr.2022.20>
- Eftaxias A, Diamantis V, Michailidis C, Stamatelatou K and Aivasidis A (2021). The role of emulsification as pre-treatment on the anaerobic digestion of oleic acid: process performance, modeling, and sludge metabolic properties. *Biomass Conversion and Biorefinery*, 11(2): 251–260. <https://doi.org/10.1007/s13399-019-00600-4>
- El-Sherbiny M, Cieślak A, Szczechowiak J, Kołodziejcki P, Szulc P and Szumacher-Strabel M (2016). Effect of nanoemulsified oils addition on rumen fermentation and fatty acid proportion in a rumen simulation technique. *Journal of Animal and Feed Sciences*, 25(2): 116–124. <https://doi.org/10.22358/jafs/65571/2016>
- Fassah DM and Khotijah L (2016). Vitamin-E supplementation on rich poly-unsaturated fatty acid ration to blood profile of lactation ewes [Pengaruh vitamin-E dalam ransum kaya asam lemak tidak jenuh terhadap profil darah induk domba laktasi]. *Jurnal Veteriner*, 17(3): 430–439. <https://doi.org/10.19087/jveteriner.2016.17.3.430>

- Geng Y, Lin G, Zhang K, Dai B, Weng Y, Wu J, et al. (2020). Effects of docosahexaenoic acid-enriched algae supplementation on growth performance, slaughter performance, muscle quality and fatty acid composition in male kids of dairy goats. *Chinese Journal of Animal Nutrition*, 32(4): 1734–1744. <https://doi.org/10.3969/j.issn.1006-267x.2020.04.032>
- Lamothe S, Desroches V and Britten M (2019). Effect of milk proteins and food-grade surfactants on oxidation of linseed oil-in-water emulsions during in vitro digestion. *Food Chemistry*, 294: 130–137. <https://doi.org/10.1016/j.foodchem.2019.04.107>
- Li H, Pan Y, Yang Z, Rao J and Chen B (2022). Modification of β -lactoglobulin by phenolic conjugations: Protein structural changes and physicochemical stabilities of stripped hemp oil-in-water emulsions stabilized by the conjugates. *Food Hydrocolloids*, 128:107578. <https://doi.org/10.1016/j.foodhyd.2022.107578>
- Liao Z, Li Q, Chen J, Du L, Wang J and Fang X (2021). Research progress on the functional activity of flaxseed oil components. *Modern Food Science and Technology*, 37(11): 379–389. <https://doi.org/10.13982/j.mfst.1673-9078.2021.11.0005>
- López-Martínez A and Rocha-Urbe A (2018). Antioxidant hydrophobicity and emulsifier type influences the partitioning of antioxidants in the interface improving oxidative stability in O/W emulsions rich in n-3 fatty acids. *European Journal of Lipid Science and Technology*, 120(1): 1700277. <https://doi.org/10.1002/ejlt.201700277>
- Losada-Barreiro S, Bravo-Díaz C and Paiva-Martins F (2020). Why encapsulate antioxidants in emulsion-based systems, where they are located, and how location affects their efficiency. In: *Emulsion-based Encapsulation of Antioxidants*. Abouzadeh MA, editor. Springer Nature, Cham. https://doi.org/10.1007/978-3-030-62052-3_1
- McDonald P, Edwards RA, Greenhalgh JD, Morgan CA, Sinclair LA and Wilkinson RG (2022). *Animal Nutrition*. 8th Edition. Pearson Education Limited, Harlow.
- Mickdam E, Khaosa-ard R, Metzler-Zebeli BU, Klevenhusen F, Chizzola R and Zebeli Q (2016). Rumen microbial abundance and fermentation profile during severe subacute ruminal acidosis and its modulation by plant derived alkaloids in vitro. *Anaerobe*, 39: 4–13. <https://doi.org/10.1016/j.anaerobe.2016.02.002>
- Mirzaei-Alamouti H, Abdollahi A, Rahimi H, Moradi S, Vazirigohar M and Aschenbach JR. (2021). Effects of dietary oil sources (sunflower and fish) on fermentation characteristics, epithelial gene expression and microbial community in the rumen of lambs fed a high-concentrate diet. *Archives of Animal Nutrition*, 75(6): 405–421. <https://doi.org/10.1080/1745039X.2021.1997539>
- Muktiani A, Widiyanti W and Pandupuspitasari NS. (2024). Supplementation of zinc palm oil soap improves feed fermentability and unsaturated fatty acid profile in rumen liquid. *Tropical Animal Science Journal*, 47(3): 371-380. <https://doi.org/10.5398/tasj.2024.47.3.371>
- Najini R and Wahdaningsih S (2024). The effect of maceration and soxhletation method on total phenolic content and antioxidant activity of jackfruit leaves (*Artocarpus heterophyllus* L.) [Pengaruh metode maserasi dan metode soxhletasi terhadap kadar fenolik total dan aktivitas antioksidan daun Nangka (*Artocarpus heterophyllus* L.)] *Jurnal Farmasi IKIFA*, 3(1): 13–25. <https://epik.ikifa.ac.id/index.php/jfi/article/view/109>
- Obando M, Papastergiadis A, Li S and De Meulenaer B (2015). Impact of lipid and protein co-oxidation on digestibility of dairy proteins in oil-in-water (O/W) emulsions. *Journal of Agricultural and Food Chemistry*, 63(44): 9820–9830. <https://doi.org/10.1021/acs.jafc.5b03563>
- Ogimoto K and Imai S (1981). *Atlas of Rumen Microbiology*. Japan Scientific Societies, Tokyo.
- Ojwang RA, Muge EK, Mbatia B and Ogoyi DO (2017). Comparative analysis of phytochemical composition and antioxidant activities of methanolic extracts of leaves, roots and bark of jackfruit (*Artocarpus heterophyllus*) from selected regions in Kenya and Uganda. *Journal of Advances in Biology & Biotechnology*, 16(1): 1–13. <https://doi.org/10.9734/JABB/2017/37355>
- Oliveira RV, Amaral L da S, Milagres CA, Santos CTB, Pelli A and Paiva AD (2021). Assessment of temperature and acid tolerance of *Bacillus subtilis* isolated from a Brazilian fruit juice-added soy beverage. *Interação*, 21(2): 38–48. <https://doi.org/10.53660/inter-105-s103-p38-48>
- Park T and Yu Z (2022). Interactions between *Entodinium caudatum* and an amino acid-fermenting bacterial consortium: fermentation characteristics and protozoal population in vitro. *Journal of Animal Science and Technology*, 65(2): 387. <https://doi.org/10.5187/jast.2022.e111>
- Pernin A, Bosc V, Soto P, Le Roux E and Maillard M (2019). Lipid oxidation in oil-in-water emulsions rich in omega-3: effect of aqueous phase viscosity, emulsifiers, and antioxidants. *European Journal of Lipid Science and Technology*, 121(9): 1800462. <https://doi.org/10.1002/ejlt.201800462>
- Pujiawati Y, Khotijah L and Wiryawan I (2023). Screening of antioxidant activities and their bioavailability of tropical plants. *IOP Conference Series: Earth and Environmental Science*, 1182(1): 012083. <https://doi.org/10.1088/1755-1315/1182/1/012083>
- Reproto RO. (2025). Effects of lipid source supplementation on rumen microbial population dynamics and in vivo digestibility of napier grass in goats. *International Journal of Agriculture and Biosciences*, 14(3): 483–489. <https://doi.org/10.47278/journal.ijab/2025.028>
- Ribeiro MD, Cabral SLS, Fabino R, Calaça AMM, Silva CJ, Zaccaroni OF, et al. (2025). Inclusion of oils in the in vitro fermentation of elephant grass (*Pennisetum purpureum* Schum. cv. Cameroon). *Revista Brasileira de Zootecnia*, 54. <https://doi.org/10.37496/rbz5420240160>
- Riestanti LU, Despal, Retnani Y and Andarwulan N (2024). Unsaturated fat supplemented in the form of Ca-soap and prill fat in dairy cattle ration: in vitro study. *BIO Web of Conferences*, 123: 01016. <https://doi.org/10.1051/bioconf/202412301016>
- Romanić RS, Lužaić TZ and Radić BĐ (2021). Enriched sunflower oil with omega 3 fatty acids from flaxseed oil: prediction of the nutritive characteristics. *LWT*, 150: 112064. <https://doi.org/10.1016/j.lwt.2021.112064>
- Roque-Jiménez JA, Rosa-Velázquez M, Pinos-Rodríguez JM, Vicente-Martínez JG, Mendoza-Cervantes G, Flores-Primo A, et al. (2021). Role of long chain fatty acids in developmental programming in ruminants. *Animals*, 11(3): 762. <https://doi.org/10.3390/ani11030762>
- Roschel GG, Silveira TFF, Cajaiba LM and Castro IA (2019). Combination of hydrophilic or lipophilic natural compounds to improve the oxidative stability of flaxseed oil. *European Journal of Lipid Science and Technology*, 121(5). <https://doi.org/10.1002/ejlt.201800459>
- Saxena N and Mandal A (2022). Surfactants in Petroleum Industry. In: *Natural Surfactants: Application in Enhanced Oil Recovery*. Springer International Publishing, Cham. pp. 11–17. https://doi.org/10.1007/978-3-030-78548-2_3
- Shen JS, Chai Z, Song LJ, Liu JX and Wu YM (2012). Insertion depth of oral stomach tubes may affect the fermentation parameters of ruminal fluid collected in dairy cows. *Journal of Dairy Science*, 95(10): 5978–5984. <https://doi.org/10.3168/jds.2012-5499>
- Sørensen A-DM, Haahr A-M, Becker EM, Skibsted LH, Bergenstahl B, Nilsson L, et al. (2008). Interactions between iron, phenolic compounds, emulsifiers, and pH in omega-3-enriched oil-in-water emulsions. *Journal of Agricultural and Food Chemistry*, 56(5): 1740–

1750. <https://doi.org/10.1021/jf072946z>

Suharti S, Aliyah D and Suryahadi S (2019). In vitro rumen fermentation characteristics with the addition of vegetable oil calcium soaps under different buffer systems [Karakteristik fermentasi rumen in vitro dengan penambahan sabun kalsium minyak nabati pada buffer yang berbeda] Jurnal Ilmu Nutrisi dan Teknologi Pakan, 16(3): 56-64. <https://journal.ipb.ac.id/index.php/jurnalntp/article/view/24939>

Tilley JMA and Terry RA (1963). A two-stage technique for the in vitro digestion of forage crops. Grass and Forage Science, 18(2): 104-111. <https://doi.org/10.1111/j.1365-2494.1963.tb00335.x>

Wardeh MF (1981). Models for Estimating Energy and Protein Utilization for Feeds. Dissertation. Utah State University, Utah, USA. <https://doi.org/10.26076/9026-5aad>

Zahera R, Pratiwi MI, Fitri A, Koike S, Permana IG and Despal (2024). Coconut fatty acid distillate Ca-soap with different calcium sources: Effects of varied proportions of protected and unprotected fat supplementation in dairy rations. Dairy, 5(3): 542-554. <https://doi.org/10.3390/dairy5030041>

Publisher's note: Scienceline Publication Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2026